

UNIVERSIDADE ESTADUAL DO MARANHÃO
CENTRO DE CIÊNCIAS AGRÁRIAS
PROGRAMA DE PÓS-GRADUAÇÃO EM AGROECOLOGIA
DOUTORADO EM AGROECOLOGIA

MARCELO MARINHO VIANA

**PRODUTIVIDADE E EFICIÊNCIA DO USO DO NITROGÊNIO NA CULTURA
DO MILHO INOCULADO COM *Azospirillum brasilense* E ADUBADO COM
MOLIBDÊNIO**

SÃO LUÍS

2021

MARCELO MARINHO VIANA
Engenheiro Agrônomo

**PRODUTIVIDADE E EFICIÊNCIA DO USO DO NITROGÊNIO NA CULTURA
DO MILHO INOCULADO COM *Azospirillum brasilense* E ADUBADO COM
MOLIBDÊNIO**

Tese apresentada ao Programa de Pós-Graduação
em Agroecologia na Universidade Estadual do
Maranhão, para obtenção do título de Doutor em
Agroecologia.

Orientador: Prof. Dr. Heder Braun

SÃO LUÍS

2021

Viana, Marcelo Marinho.

Produtividade e eficiência do uso de nitrogênio na cultura do milho inoculado com *Azospirillum brasiliense* e adubado com molibdênio / Marcelo Marinho Viana. – São Luís, 2021.

105 f

Tese (Doutorado) - Programa de Pós-Graduação em Agroecologia, Universidade Estadual do Maranhão, 2021.

Orientador: Prof. Dr. Heder Braun.

1.Trópico úmido. 2.Bactérias diazotróficas. 3.Nutrição de plantas. I.Título.

CDU:633.15-184.7

Elaborado por Giselle Frazão Tavares - CRB 13/665

MARCELO MARINHO VIANA

**PRODUTIVIDADE E EFICIÊNCIA DO USO DO NITROGÊNIO NA CULTURA
DO MILHO INOCULADO COM *Azospirillum brasilense* E ADUBADO COM
MOLIBDÊNIO**

Aprovado em: 30 de dezembro de 2021

BANCA EXAMINADORA




Dr. Heder Braun (Orientador)
Universidade Estadual do Maranhão



Dr. Marlon Gomes da Costa
Instituto Federal de Educação, Ciência e Tecnologia do Maranhão



Dra. Katia Pereira Coelho
Universidade Federal Rural do Rio de Janeiro



Dra. Camila Nobre Pinheiro
Universidade Estadual do Maranhão



Dr. João Batista Zonta
Empresa Brasileira de Pesquisa Agropecuária

Dedico

Aos meus pais,
José de Jesus Neves Viana e Maria José Marinho Viana,
por todo amor e dedicação incondicional aos filhos,
pelo incentivo e orientação.

AGRADECIMENTOS

Primeiramente a **Deus**, pela sua bondade e misericórdia, por me conceder saúde e proteção divina. Obrigado por mais esta conquista, foram longos 11 anos para chegar até aqui.

Aos meus pais, **José de Jesus Neves Viana** e **Maria José Marinho Viana**, por todos os ensinamentos, carinho, zelo, amor, companheirismo, incentivo, pela força que me deram durante toda a caminhada, enfim, por tudo. Nada teria sido possível sem vocês!

Agradeço a todos os meus familiares, mas em especial aos meus irmãos Leonardo de Jesus Marinho Viana, Higo Bruno Brito Marinho e Ana Leticia Garcia Viana, pelo apoio e credibilidade que sempre me impulsionaram a ir mais longe, a fazer mais, a ser melhor. Sintam-se parte de mais esta conquista.

À minha esposa Bruna Campos Sena e meu filho João Gabriel Sena Viana, pelo companheirismo, força, incentivo, paciência e todo amor, ao qual me capacita a ser uma pessoa melhor e enfrentar os percalços caminhada que se chama vida.

À minha segunda mãe Maria José Gomes da Silva, por todo amor, carinho, incentivo, em todos os momentos, que possibilitaram chegar ao final desse objetivo.

À Universidade Estadual do Maranhão (UEMA), ao Programa de Pós-Graduação em Agroecologia, Bangor University e CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior) pela oportunidade de realização deste sonho em minha vida e alastrar meus conhecimentos

Ao meu amigo e orientador Dr. Heder Braun, sou grato por tudo que aprendi e vivi com ele, pois é uma pessoa que orienta para vida com seus conselhos e ensinamentos profissionais. Agradeço pela confiança e parceria durante todo esse tempo em que estivemos juntos.

A todo corpo docente do Programa de Pós-Graduação em Agroecologia, especialmente ao professor Dr. Fabricio de Oliveira Reis e à professora Dr^a. Antônia Alice Costa Rodrigues pela colaboração e disponibilidade sempre.

Agradeço carinhosamente à Rayanne Cristine e Denise, secretárias do Programa de Pós-Graduação, por sua prontidão em servir e ajudar, sem medir esforços.

Aos pesquisadores Dra. Cristina Carvalho, Dra. Camila Nobre Pinheiro e Dra. Katia Coelho por todo incentivo e ajuda.

Aos funcionários Joãozinho e Neto, por toda ajuda com a realização do trabalho em campo, sem eles seria impossível realizar todos os experimentos, registro aqui meus sinceros agradecimentos.

A todos os amigos que diretamente ou indiretamente contribuíram para a realização desse trabalho. Especialmente a Luís Carlos Ferreira Reis, Werlen Barbosa, Ester de Paiva Alves, Erivaldo Plinio, Assistone Costa, Marcela Uli, Eluado Marques, Benjamim Valentim, Henry Reyes.

Aos meus irmãos de pai científico: Karen Alessandra, Lincon Matheus, Werlen Barbosa, Ester de Paiva Alves, Erivaldo Plinio, Danilo Sodré, Vanessa Lira e Fernando pelos incontáveis dias de trabalho, alegria, discussões e parceria.

À FAPEMA pela concessão da bolsa de estudos e pelo apoio financeiro na execução do projeto.

Obrigado a todos!

“Tudo posso naquele que me fortalece”

Filipenses 4, 13

LISTA DE ILUSTRAÇÕES

CAPÍTULO II

Pg.

Figure 1. Rainfall (mm) and maximum and minimum temperatures obtained from the data base National Institute of Meteorology of Brazil (INMET) during the corn cultivation (all seasons) in the period from January 2018 to July 2019. 44

CAPÍTULO III

Figure 1. Rainfall (mm) and maximum and minimum temperatures obtained from the data base National Institute of Meteorology of Brazil (INMET) during the corn cultivation (all seasons) in the period from January 2018 to July 2019. 69

LISTA DE TABELA

CAPÍTULO II	Pg.
Table 1. Treatment arrangement of all seasons 2018/2019 with <i>Azospirillum brasilense</i> (seeds and leaf), nitrogen and molybdenum rates.	45
Table 2. Mean value (\pm SD) of grain yield, 100-seed weight, harvest index, nitrogen use efficiency (NUE) and leaf chlorophyll index (LCI), experiments 1,2 and 3 in the maize under field conditions in research farm, Maranhão State University, Maranhão State, Brazil.	61
Table 3. Mean value (\pm SD) of plant height, stem diameter, insertion height of the first ear (IHFE) and shoot dry matter, experiments 1,2 and 3.	62
CAPÍTULO III	
Table 1. Treatment arrangement of all seasons 2018/2019 with <i>Azospirillum brasilense</i> (seeds and leaf), nitrogen and molybdenum rates.	70
Table 2. Effect of treatments evaluated at 20 days after topdressing, mean value (\pm SD) of CO ₂ assimilation (A, $\mu\text{mol CO}_2\cdot\text{m}^{-2}\cdot\text{s}^{-1}$), stomatal conductance ($\text{gs mol m}^{-2}\cdot\text{s}^{-1}$), intercellular CO ₂ concentration (C _i , $\mu\text{mol CO}_2/\text{mol}^{-1}$) and leaf chlorophyll index (LCI). Experiments 1,2 and 3.	85
Table 3. Effect of treatments evaluated at 20 days after topdressing, mean value (\pm SD) of photosynthetic nitrogen use efficiency (PNUE, $\mu\text{mol CO}_2\text{ g}^{-1}\cdot\text{N}\cdot\text{s}^{-1}$). Experiments 1,2 and 3.	86
Table 4. Effect of treatments evaluated at 40 days after topdressing, mean value (\pm SD) of CO ₂ assimilation (A, $\mu\text{mol CO}_2\cdot\text{m}^{-2}\cdot\text{s}^{-1}$), stomatal conductance ($\text{gs mol m}^{-2}\cdot\text{s}^{-1}$), intercellular CO ₂ concentration (C _i , $\mu\text{mol CO}_2/\text{mol}^{-1}$) and leaf chlorophyll index (LCI). Experiments 1,2 and 3.	87
Table 5. Effect of treatments evaluated at 40 days after topdressing, mean value (\pm SD) of photosynthetic nitrogen use efficiency (PNUE, $\mu\text{mol CO}_2\text{ g}^{-1}\cdot\text{N}\cdot\text{s}^{-1}$). Experiments 1,2 and 3.	88

RESUMO

O nitrogênio (N) é um dos nutrientes mais importantes e exigidos pela cultura do milho e o que mais onera a produção agrícola, pois na maioria dos solos agricultáveis este elemento não está disponível em níveis necessários para atingir altas produtividades. Uma alternativa é a utilização de produtos biológicos (bactérias do gênero *Azospirillum*) e o micronutriente molibdênio (Mo), que também é importante, pois esse micronutriente interfere diretamente no metabolismo do N. O objetivo desse trabalho foi investigar o efeito da inoculação com *Azospirillum brasilense* associada a aplicação de Mo e N no desenvolvimento produtivo e na eficiência do uso do N e eficiência fotossintética do N em plantas de milho cultivada em solo da região trópico úmido maranhense. Três experimentos de campo foram conduzidos no município de São Luís (2° 30' S, 44° 18' O, 24 metros acima do nível do mar), Estado do Maranhão, Brasil. O delineamento experimental foi em blocos casualizados com quatro repetições e oito tratamentos. Quando o efeito do tratamento foi significativo, sete contrastes ortogonais foram analisados. Os tratamentos foram uma dose de N 140 kg ha⁻¹, uma dose de Mo 90 g ha⁻¹ e métodos de inoculação de *A. brasilense* (semente e foliar). As variáveis dependentes analisadas foram: índice de clorofila foliar, altura da planta, diâmetro do caule, altura de inserção da primeira espiga, peso de 100 sementes, produtividade de grãos, índice de colheita, matéria seca da parte aérea e eficiência de uso de nitrogênio. Avaliamos também características fisiológicas no início do estágio de florescimento (aproximadamente 20 dias após a adubação de cobertura) e do grão leitoso (aproximadamente 40 dias após a adubação de cobertura) as características foram: assimilação fotossintética de CO₂, condutância estomática, concentração intercelular de CO₂, índice de clorofila foliar e a eficiência do uso de N fotossintético. A inoculação com *A. brasilense*, nitrogênio e Mo mostrou-se potencialmente utilizável pois os resultados mostraram que a combinação (*A. brasilense*, Mo e N) aumentou 5,1% da eficiência do uso do nitrogênio e 12,68% da eficiência do uso de N fotossintético. No entanto, não houve aumento na produtividade de grãos. Por outro lado, Mo e N combinados aumentaram 17,8% a produtividade de grãos quando comparadas as plantas de milho inoculadas apenas com *A. brasilense*. Diante desses resultados é necessário propor novas pesquisas com o papel de elucidar melhor o papel do Mo na ligação com bactérias promotoras de crescimento e seus efeitos na rizosfera e na produtividade de grãos.

Palavras-chave: Tropicó úmido, bactérias diazotróficas, nutrição de plantas.

ABSTRACT

Nitrogen (N) is one of the most important nutrients required by the maize crop and the most burdensome on agricultural production since in most arable soils this element is not available at the levels necessary to achieve high yields. An alternative is the use of biological products (bacteria of the genus *Azospirillum*) and the micronutrient molybdenum (Mo), which is also important, as this micronutrient directly interferes with N metabolism. The objective of this work was to investigate the effect of inoculation with *A. brasilense* associated with the application of Mo and N on the productive development and on the N use efficiency and N photosynthetic efficiency in maize plants grown in soil from the humid tropic region of Maranhão. Three field experiments were carried out in the municipality of São Luís (2° 30' S, 44° 18' W, 24 meters above sea level), State of Maranhão, Brazil. The experimental design was in randomized blocks with four replications and eight treatments. When the treatment effect was significant, seven orthogonal contrasts were analyzed. The treatments were a dose of N 140 kg ha⁻¹, a dose of Mo 90 g ha⁻¹ and methods of inoculation of *A. brasilense* (seed and leaf). The dependent variables analysed were: leaf chlorophyll index, plant height, stem diameter, height of insertion of the first ear, weight of 100 seeds, grain yield, harvest index, shoot dry matter and nitrogen use efficiency. We also evaluated physiological characteristics at the beginning of the flowering stage (approximately 20 days after top dressing) and milky grain (approximately 40 days after top dressing). The characteristics were: photosynthetic CO₂ assimilation, stomatal conductance and intercellular concentration of CO₂, leaf chlorophyll index and photosynthetic N use efficiency. Inoculation with *A. brasilense*, nitrogen and Mo proved to be potentially usable as the results showed that the combination (*A. brasilense*, Mo and N) increased 5.1% of the nitrogen use efficiency and 12.68% of the photosynthetic nitrogen use efficiency. However, there was no increase in grain yield. On the other hand, Mo and N combined increased grain yield by 17.8% when compared to corn plants inoculated only with *A. brasilense*. Given these results, it is necessary to propose further research with the role of better elucidating the role of Mo in the link with growth-promoting bacteria and its effects on the rhizosphere and on grain yield.

Keywords: Humid tropic, diazotrophic bacteria, plant nutrition.

SUMÁRIO

CAPÍTULO I	15
1. INTRODUÇÃO GERAL	16
2. REVISÃO BIBLIOGRÁFICA	19
2.1. <i>Cultura do milho</i>	19
2.2. <i>Nitrogênio</i>	20
2.3. <i>Eficiência do Uso do Nitrogênio (EUN)</i>	21
2.4. <i>Fixação Biológica do Nitrogênio (FBN)</i>	23
2.5. <i>Azospirillum brasilense</i>	24
2.6. <i>Molibdênio</i>	26
REFERÊNCIAS BIBLIOGRÁFICAS	28
The effects of <i>Azospirillum brasilense</i>, nitrogen and molybdenum on the grain yield and nitrogen use efficiency on maize plants grown in the sub-humid tropics of Brazil.	39
ABSTRACT	41
1. INTRODUCTION	41
2. MATERIALS E METHODS	43
2.1. <i>Field site description</i>	43
2.1. <i>Experimental design, treatment, and field management</i>	45
2.3. <i>Measurements collected</i>	47
2.4. <i>Statistical analyses</i>	48
3. RESULTS	49
4. DISCUSSION	51
5. CONCLUSIONS	54
Acknowledgements	54
References	55
Application methods of <i>Azospirillum brasilense</i> associated with nitrogen and molybdenum on ecophysiological parameters on maize plants cultivated in the sub-humid tropical regions of Brazil	63
ABSTRACT	65
1. INTRODUCTION	65
2. MATERIALS E METHODS	68

<i>2.1. Field site description</i>	68
<i>2.2. Experimental design, treatment, and field management</i>	70
<i>2.3. Measurements collected</i>	72
<i>2.4. Statistical analyses</i>	72
3. RESULTS	73
6. DISCUSSION	76
7. CONCLUSIONS	78
Acknowledgements	78
References	79
CONSIDERAÇÕES FINAIS	89
ANEXOS	90

REFERENCIAL TEÓRICO

CAPÍTULO I

1. INTRODUÇÃO GERAL

A agricultura é uma atividade indispensável para o mundo porque desempenha um papel importante no fornecimento de alimentos para a população mundial. Por outro lado, a pesquisa agrícola tem um problema real e urgente a ser resolvido: produzir mais alimentos em um espaço menor, ou seja, aumentar a produtividade das lavouras. O foco é usar um método chamado "land sparing – efeito poupa-terra": para produzir mais alimentos em um espaço menor. Isso visa não expandir a área cultivável, mas intensificar a produção e fortalecer a proteção ambiental alocando mais espaço para fitorremediação (técnica de descontaminação de solo e água, através da utilização de plantas). Atualmente a corrida pela não expansão de novas áreas agrícolas, tem levado pesquisadores a se esforçarem para aumentar a produtividade das lavouras, com foco na quantidade e a qualidade dos produtos, a viabilização dos custos de investimento e a concretização final de planos ecologicamente corretos e socialmente justos. Nesse campo, podemos focar na cultura do milho (*Zea mays* L.), por ser de grande importância econômica e alimentar.

O milho é uma cultura de importância global em virtude da sua diversidade de utilização, extensão da área cultivada e elevada capacidade produtiva. Os maiores produtores mundiais são os Estados Unidos, Canadá e o Brasil. Na safra de 2019/2020 o Brasil obteve produtividade média de 5,531 Mg ha⁻¹ (CONAB, 2020).

É sabido que as lavouras de milho são altamente dependentes de um bom manejo de fertilizantes de nitrogênio. Embora o manejo da fertilização das lavouras de milho tenha melhorado, o nitrogênio (N) ainda é um dos principais fatores que levam à baixa produtividade das lavouras de milho (USDA, 2017). O N é o nutriente mais necessário no ciclo produtivo da lavoura de milho, e sua deficiência pode causar perdas significativas de produtividade. Os fertilizantes nitrogenados respondem por grande parte do custo de fertilização das lavouras de milho, que pode variar dependendo da produtividade necessária (DARTORA *et al.*, 2013). A maioria dos solos nos trópicos úmidos é deficiente em nitrogênio, apenas 30% a 40% do nitrogênio é usado pelas lavouras e 60% a 70% do nitrogênio é perdido por lixiviação, volatilização e desnitrificação (HIREL *et al.*, 2007; KONG *et al.*, 2016).

A fixação biológica de nitrogênio (FBN) é benéfica para o suprimento de parte do N necessário à cultura do milho durante seu ciclo produtivo. Nas monocotiledôneas, a FBN ocorre por meio de bactérias diazotróficas, ao qual iremos destacar o gênero *Azospirillum*. Esses microrganismos são capazes de reduzir o N atmosférico (N₂) a amônia (NH₄⁺) pela quebra da ligação tríplice do N através da enzima nitrogenase, com alto consumo de energia na forma de

adenosina trifosfato (ATP) (SANGOI *et al.*, 2015). As bactérias diazotróficas também podem atuar no crescimento vegetal, produção de hormônios (auxinas, citocininas, giberelinas, etileno) e atuam como agentes de controle biológico de patógenos (CORREA *et al.*, 2008). Em geral, as bactérias diazotróficas geram benefícios ao desenvolvimento das plantas pela combinação de todos esses mecanismos listados acima (DOBBELAERE *et al.*, 2003). No entanto, a inconsistência dos resultados de pesquisas inviabiliza inferências sobre as possíveis interações das bactérias diazotróficas e a disponibilidade de N mineral no que diz respeito à resposta da planta (BREDA *et al.* 2020), pois o nitrato (NO_3^-) também é um modulador da arquitetura da raiz e pode atuar em um papel sinérgico na melhoria da exploração do solo pelas plantas (FORDE 2014). Nesse sentido, a genética das plantas e o ambiente afetam o metabolismo das bactérias diazotróficas e, conseqüentemente, a resposta de promoção do crescimento das plantas. Os estudos genéticos são fundamentais para descrever e revelar as etapas e funcionalidades das bactérias diazotróficas e seus efeitos nas monocotiledôneas.

Diante desse cenário, uma alternativa para redução da dependência de fertilizantes nitrogenados nas lavouras de milho, é a utilização consorciada de bactérias diazotróficas do gênero *Azospirillum* e o micronutriente molibdênio (Mo). Essa associação tem como objetivo, aumentar o rendimento produtivo da lavoura de milho e manter o balanço nutricional adequado.

O Mo é um micronutriente exigido pelas plantas em pequenas quantidades. No entanto, sua deficiência é tão prejudicial quanto a falta de um macronutriente (N, K e P), principalmente pois afeta diretamente o metabolismo do nitrogênio (TAIZ; ZEIGER, 2017). A utilização de N pelas plantas de milho pode ser potencializada com o uso de Mo, uma vez que, este micronutriente é constituinte de enzimas que permitem a assimilação de N via fixação biológica por microrganismos diazotróficos (nitrogenase) e/ou N sintético (nitrato redutase) (PICAZEVICZ *et al.*, 2017).

As plantas absorvem Mo na forma de molibdato (MoO_4^{2-}) (VALENTINI *et al.*, 2005). Embora a quantidade necessária seja pequena, o teor nos tecidos da planta é geralmente inferior a $1,0 \text{ mg kg}^{-1}$ de matéria seca. (MENGEL; KIRKBY, 1987). Estudos constataram que o uso de Mo com *Azospirillum brasilense* (GANAPATHY; SAVALGI, 2006) e fertilizantes nitrogenados sintéticos (VALENTINI *et al.*, 2005) pode aumentar o crescimento e a produtividade das plantas de milho. No entanto, o efeito da pulverização de Mo na produtividade do milho é diferente, dependendo da localização e do tipo de solo.

Diante dos relatos apresentados, a comunidade científica possui um déficit de informações sobre o uso de bactérias diazotróficas (*Azospirillum brasilense*) associadas a adubação com o micronutriente Mo e seus efeitos em plantas de milho em solo da região do

tropico úmido maranhense, o que pode contribuir para a redução parcial de uso do N sintético aplicado em cobertura e potencializar os benefícios dessa técnica com o consequente aumento da produtividade.

2. REVISÃO BIBLIOGRÁFICA

2.1. Cultura do milho

O milho (*Zea mays* L.), é uma espécie, pertencente à família das Poáceas (antiga família das gramíneas), possivelmente tem origem americana, foi encontrado em pequenas ilhas próximas ao litoral mexicano. Para alguns membros da comunidade científica, o milho é originário do Teosinte (*Zea mexicana* L.), uma gramínea anual originária do México e da Guatemala, por meio de seleção feita pelo homem, outros membros defendem a hipótese de que o milho e o Teosinte diferenciam-se a mais tempo de um ancestral comum (SOUZA, 2017).

Após o descobrimento das Américas, o milho foi levado para o continente europeu, onde plantas foram cultivadas em jardins até seu valor alimentício tornar-se conhecido (OKUMURA *et al.*, 2011). Na contemporaneidade, devido à sua alta capacidade de adaptação a diversas condições de cultivo, e em virtude da grande variedade de genótipos existentes, a cultura do milho está presente em todos os continentes e sua produção mundial só perde para a cultura do trigo e a do arroz (SILVA *et al.*, 2014).

A cultura do milho, possui papel incontestável na economia mundial e brasileira devido a sua posição entre as espécies agrícolas com maior área de cultivo (MÔRO; FRITSCHÉ, 2015). No Brasil, os maiores produtores de milho na safra de 2019/2020 foram os estados de Mato Grosso, Paraná, Goiás, Mato Grosso do Sul e Minas Gerais. O estado do Maranhão participou com uma produção de aproximadamente 977,3 mil toneladas (CONAB, 2020).

Nas condições de exploração do milho no Brasil, grande parte das cultivares possui ciclo de produção entre 105 e 180 dias, período compreendido entre a semeadura e a colheita (MOREIRA, 2015). Segundo Magalhães *et al.* (2006) a cultura do milho, para expressar seu potencial produtivo, necessita em média de 600 mm de precipitação e temperaturas entre 25 e 30 °C.

A importância econômica do milho grão é caracterizada pelas diversas formas de sua utilização, que vai desde a alimentação animal, como forragem conservada para o período da seca e na fabricação de farelos até a indústria química, como matéria prima de mais de 500 produtos e a indústria alimentícia, como amido, farinhas e óleo (OKUMURA *et al.*, 2011).

Há vários fatores que limitam a produtividade do milho, como por exemplo, doenças, pragas, déficit hídrico, disponibilidade e assimilação de nutrientes. Em relação à fertilidade, a cultura do milho é muito exigente em fertilizantes, principalmente nitrogenados (FERNANDES *et al.*, 2008). Neste sentido, estudos devem ser intensificados com o intuito de

desenvolver técnicas alternativas e promissoras para a eficácia da incorporação do nitrogênio nas plantas de milho, o que refletirá em estratégia econômica e de menor impacto ambiental.

2.2. Nitrogênio

O nitrogênio (N) é um nutriente essencial absorvido pelas raízes, convertido em aminoácidos para compor diversas moléculas nas raízes e na parte aérea das plantas durante todo o período de crescimento (WANG; XING, 2017). É importante, sobretudo nos estádios iniciais do desenvolvimento vegetal, quando sua disponibilidade se relaciona diretamente com as maiores eficiências de utilização pelas plantas. Nas plantas de milho é importante na fase de aproximadamente quatro folhas, quando se define o potencial produtivo (RITCHIE *et al.*, 2003).

O N possui papel fundamental no metabolismo vegetal por participar diretamente na biossíntese de proteínas e clorofilas (ANDRADE *et al.*, 2003). É constituinte de proteínas, enzimas, coenzimas, ácidos nucleicos, fitocromos e integra a molécula da clorofila. Além disso, afeta as taxas de iniciação e expansão foliar, o tamanho final e a intensidade de senescência das folhas, o desenvolvimento da área foliar e a taxa de fotossíntese, o crescimento radicular, o rendimento biológico, o tamanho e número de espigas, a massa de grãos e índice de espiga, a altura de planta, o comprimento da espiga, o diâmetro de colmo, a inserção de espiga, o número de plantas acamadas e quebradas e a qualidade de grãos (SILVA, 2014). A maioria dos fertilizantes nitrogenados empregados na cultura milho são hidrossolúveis, e rapidamente liberam para o solo NO_3^- e NH_4^+ , sendo desta forma prontamente assimiláveis pela planta. Assim, as plantas de milho, por remover grandes quantidades de nitrogênio, requer o uso de adubação nitrogenada em cobertura para complementar a quantidade suprida pelo solo. O N é predominantemente derivado de fertilizantes, fixação biológica de N_2 , mineralização do N orgânico de esterco animal, resíduos de culturas e matéria orgânica do solo (SOUZA, 2017), no entanto, por sua alta mobilidade, o N está susceptível a perdas e pode ocasionar riscos ambientais (MENDES, 2016).

Em grande parte dos solos brasileiros, a quantidade de N é insuficiente, fazendo-se necessário o fornecimento externo do nutriente em concentração adequada para garantir o crescimento, desenvolvimento e a produtividade das plantas de milho (BELARMINO *et al.*, 2003), e isso, deve ser feito respeitando-se os limites do solo, para evitar a degradação. Nesse contexto, o manejo da adubação nitrogenada é realizado com o intuito de garantir boa produtividade, em função da dinâmica do N, grandes quantidades desse nutriente são

adicionadas ao solo, o que pode levar a perdas e degradação do ambiente. E tendo em vista a crescente demanda por fertilizantes nitrogenados e a preocupação com as possíveis perdas e contaminação do ambiente (FERNANDES; LIBARDI, 2007), faz-se necessária a aplicação de N na forma parcelada em cobertura (YAMADA; ABDALA, 2000), como também, a investigação de alternativas para suprimento de N via fixação biológica ou fertilizantes alternativos. Isso reduzirá as perdas do nutriente e aumentará sua eficiência de uso (BOTREL *et al.*, 1999).

Com esse conhecimento, surge a necessidade de buscar alternativas para diminuir as perdas através do parcelamento da adubação de cobertura, assim como buscar alternativas para suplementar com N os plantios. Uma opção é favorecer a fixação biológica de N, essa técnica pode incrementar o rendimento da cultura do milho sem prejuízos aos recursos naturais (BASI, 2013).

2.3. Eficiência do Uso do Nitrogênio (EUN)

Nas últimas décadas, os esforços têm sido direcionados no sentido de otimizar a eficiência de utilização de nutrientes pelas plantas, visando reduzir os custos de produção, evitar a degradação dos recursos ambientais e aumentar o rendimento das culturas (KOLCHINSKI; SCHUCH, 2003). Nesse sentido, existem diversos caminhos possíveis para aumentar a eficiência do uso do nitrogênio (EUN). Um dos caminhos mais simples é a redução nas doses de adubos nitrogenados para níveis que sejam produtivos e seguros ambientalmente (FERNÁNDEZ *et al.* 1998).

O uso de fertilizantes em culturas de grãos e fibras também é importante na manutenção das reservas de N do solo. Alta produtividade com doses baixas de N, normalmente significa que a quantidade de N exportada com a colheita é maior do que a adicionada, o que contribui para o empobrecimento do solo (ALVES *et al.*, 2006). Fernandes *et al.* (2005) estudando doses de N em seis cultivares de milho e a eficiência de uso desse nutriente pela cultura, em região de cerrado, verificaram que a eficiência do uso de nitrogênio de todos os híbridos testados diminuiu com o aumento da dose de N aplicada. Além disso, observaram ainda que as doses de N influenciaram principalmente a massa de 100 grãos e a produtividade de grãos. Dessa forma, nos estudos sobre a dinâmica do N no sistema solo-planta, muitas vezes, é difícil quantificar a origem deste nutriente (SCIVITTARO *et al.*, 2000).

A eficiência nutricional, pode ser definida como a quantidade de matéria seca ou grãos produzidos por unidade de nutriente aplicado (FAGERIA, 1998). A eficiência da utilização do

nitrogênio adicionado ao solo, por sua vez, se refere ao grau de recuperação desse elemento pelas plantas, considerando as perdas que geralmente ocorrem (BREDEMEIER; MUNDSTOCK, 2000). A eficiência nutricional depende de vários processos fisiológicos, tais como absorção, assimilação e retranslocação do nitrogênio pela planta (MOLL *et al.*, 1982; CARVALHO, 2011) e pode ser aumentada com a adoção de práticas de manejo apropriadas.

Especificamente em milho, Moll *et al.* (1982) definiram a eficiência de uso do N (EUN) como a massa de grãos dividida pela massa de N aplicado no solo (Gw/Ns), ambas expressas na mesma unidade, como por exemplo, gramas por planta. Entretanto, a produtividade de grãos também se destaca como bom parâmetro da eficiência de utilização do N (CARVALHO, 2011). Para Fageria (1998) ao avaliar experimentos de campo, a produção de grãos foi o melhor parâmetro para avaliação da eficiência nutricional em culturas anuais.

Os principais componentes de avaliação da EUN são a eficiência de absorção do N (EAN), a eficiência de utilização do N (EUtN) e a eficiência de remobilização do N (ERN) (LE GOUIS *et al.*, 2000). A EAN representa a capacidade das plantas em absorver o N disponível do solo. A EUtN é definida como a relação entre o rendimento da cultura e o N total absorvido pela planta (N nos grãos + N na biomassa), ou seja, essa medida indica o rendimento de grãos obtido para cada unidade de N absorvido pela planta. Em adição, a eficiência de remobilização do N (ERN) representa a capacidade das plantas em translocar o N após a antese da parte vegetativa para os grãos. Cultivares com uma maior ERN tendem a acelerar a senescência e aumentar os níveis de N nos grãos (GAJU *et al.*, 2014).

Segundo Fidelis *et al.* (2007), a identificação de genótipos capazes de absorver e utilizar o nitrogênio de forma eficiente é um dos caminhos para aumentar a EUN na cultura do milho, incrementar a produção, minimizar as perdas e reduzir a contaminação do meio ambiente. Entretanto, a melhor eficiência nutricional é aquela obtida sob nível de nutriente adequado em que a produtividade máxima é obtida, visto que a eficiência nutricional diminui com níveis crescentes de um nutriente, devido ao suprimento desse nutriente exceder as necessidades da cultura (FAGERIA, 1998). Fernandes *et al.* (2005) e Farinelli; Lemos (2010), relataram que a eficiência do uso de nitrogênio em todos os híbridos testados nas pesquisas, diminuiu com o aumento da dose de N aplicada.

Nesse contexto, o conhecimento da relação entre os caracteres envolvidos na eficiência nutricional e o uso racional da adubação nitrogenada é fundamental, não apenas para aumentar a eficiência de recuperação do nitrogênio, mas também para aumentar a produtividade da cultura e diminuir o custo de produção (FAGERIA *et al.*, 2007).

2.4. Fixação Biológica do Nitrogênio (FBN)

O nitrogênio ainda que seja o gás mais abundante na atmosfera, não é prontamente assimilado pelas plantas. A forma que as plantas assimilam o N difere entre as espécies vegetais, as quais absorvem principalmente as formas inorgânicas deste nutriente, o nitrato ou o amônio (WILLIAMS; MILLER, 2001; FAGERIA *et al.* 2003; SOUZA; FERNANDES, 2006).

O nitrogênio gasoso (N_2) compõe 78% da atmosfera terrestre. A fixação de nitrogênio requer a quebra da ligação tripla covalente de excepcional estabilidade do N_2 . Contudo, os gases atmosféricos também se difundem para o espaço poroso do solo e o N_2 pode ser aproveitado por alguns microrganismos, principalmente bactérias que ali habitam, graças à ação da enzima dinitrogenase, que tem a capacidade de romper a tripla ligação do N_2 e reduzi-lo a NH_3 , a mesma forma obtida no processo industrial (HUNGRIA *et al.*, 2011; TAIZ; ZEIGER, 2017).

A fixação biológica de nitrogênio (FBN) é um processo de transformação do N_2 na forma inorgânica combinada NH_3 . A FBN envolve uma sucessão de processos que começam com a adaptação da bactéria à planta e culminam na fixação do N_2 atmosférico (FAGAN, *et al.*, 2007)

Todo o sistema de fixação biológica de nitrogênio é coordenado pelo complexo enzima nitrogenase, que é formado por duas unidades proteicas, a ferro-proteína (Fe-proteína) e a molibdênio-ferro-proteína (MoFe-proteína) que são responsáveis pela fixação de nitrogênio (BURRIS, 1999; MYLONA *et al.*, 1995; TAÍZ; ZIEGER, 2017). Para que ocorra a FBN é necessário que a nitrogenase esteja em condições anaeróbicas. Na reação de redução do N_2 a nitrogenase é auxiliada por uma enzima transportadora de elétrons, a ferredoxina, originária do fotossistema I da fase fotoquímica da fotossíntese (BURRIS, 1999; TAIZ; ZIEGER, 2017).

A capacidade de reduzir o nitrogênio atmosférico a amônia, está restrita a um pequeno grupo de microrganismos denominados diazotróficos, ou fixadores de N_2 (NOVAKOWISKI *et al.*, 2011). Pesquisas sobre as bactérias diazotróficas tiveram início no Brasil há mais de 40 anos. Essas pesquisas foram realizadas pela renomada pesquisadora Johanna Döbereiner e seus colaboradores (MOREIRA *et al.* 2010).

Esses microrganismos estão distribuídos em diversos grupos filogenéticos e habitam vários ecossistemas em vida livre, em simbiose com leguminosas como o feijoeiro e o feijão caupi, ou endofiticamente em raízes ou parte aérea de poaceae como milho, bem como de

espécies forrageiras como *Brachiaria* ssp., *Paspallum notatum* F. entre outras (WEBER *et al.*, 2000; MOREIRA *et al.*, 2010).

Dentre as diazotróficas, bactérias do gênero *Azospirillum* associam-se à rizosfera da planta de milho e podem contribuir com a nutrição nitrogenada da cultura (FIGUEIREDO *et al.*, 2009). Pesquisa realizada por Novakowiski *et al.* (2011), demonstrou que as bactérias do gênero *Azospirillum* podem promover o crescimento vegetal através da produção de fitoreguladores e sideróforos ou por aumentar a disponibilidade de fósforo.

No processo de FBN em gramíneas, somente uma parte do nitrogênio fixado diretamente para a planta associada é secretado para suprir parcialmente suas necessidades. Já em leguminosas, a inoculação das culturas com microrganismos, ainda que fixem nitrogênio, não conseguem suprir totalmente as necessidades das plantas em relação ao N (HUNGRIA, 2011).

O grande interesse na fixação biológica em gramíneas é devido à maior facilidade de aproveitamento de água das mesmas em relação às leguminosas, pela maior efetividade fotossintética. As gramíneas apresentam um sistema radicular fasciculado, tendo vantagens sobre o sistema pivotante das leguminosas para extrair água e nutrientes do solo; e por serem as gramíneas largamente utilizadas como alimento pelo homem. Por isso, a FBN é um processo importante para a economia em adubos nitrogenados e equilíbrio ambiental (DÖBEREINER, 1992).

2.5. *Azospirillum brasilense*

As bactérias diazotróficas não simbióticas (BDNS) atuam no desenvolvimento das plantas por meio da FBN e também pela produção e liberação de substâncias reguladoras do crescimento vegetal (SILVA; MELLONI, 2011). No grupo das bactérias diazotróficas não simbióticas (BDNS), destaca-se o gênero *Azospirillum*. Pesquisas consideram as bactérias do gênero *Azospirillum* como diazotróficas facultativas, capazes de colonizar raízes de plantas não leguminosas interna e externamente (BALDANI *et al.*, 1997). Esse gênero é caracterizado por organismos de metabolismo bastante versátil, o que confere características adaptativas, que permitem a sobrevivência em meio nutritivo rico e protetor existente na rizosfera das plantas (STEENHOUDT; VANDERLEYDEN, 2000). As BDNS podem desempenhar importante papel na sustentabilidade dos ecossistemas, uma vez que incorporam N₂ por meio da fixação biológica (DOBBELAERE *et al.*, 2003).

A fixação de nitrogênio pelas bactérias diazotróficas não simbióticas (BDNS) ocorre em ambiente natural, temperatura ambiente e seu crescimento ideal varia numa faixa de temperatura entre 28 e 41° C (ECKERT *et al.*, 2001), ocorrem em níveis bem menores de energia, consumindo os açúcares da planta, mas que são compensados pelo aporte de N fornecido ao sistema (ALVES, 2007).

As bactérias do gênero *Azospirillum* são de vida livre, rizobactérias capazes de promover o crescimento das plantas e aumentar a produtividade em muitas culturas de importância econômica. Essas bactérias podem atuar no crescimento da planta através das sínteses de hormônios, principalmente auxinas (ácido 3-indolacético), giberelinas e citocininas (FIGUEIREDO *et al.*, 2009). Ocorre também a síntese de etileno, podendo agir como solubizador de fosfato ou acelerador do processo de mineralização (PERSELLHO-CARTINEAUX *et al.*, 2003; SÁ JÚNIOR, 2012). Todo esse processo de síntese tem como consequência uma maior absorção de água e nutrientes (CORREA *et al.*, 2008) resultando em uma planta mais vigorosa e produtiva (BASHAN *et al.*, 2004; HUNGRIA, 2011).

As bactérias diazotróficas foram descobertas no início da década de 1970 pela pesquisadora da Embrapa Dr^a. Johanna Döbereiner. Essas bactérias auxiliam por diversos mecanismos na nutrição nitrogenada das culturas. Dentre esses mecanismos, destacam-se a produção de hormônios, que interferem no crescimento das plantas e podem alterar a morfologia das raízes, possibilitando a exploração de maior volume de solo (BASHAN; HOLGUIN, 1997; ZAIED *et al.*, 2003), o aumento do processo da redução assimilatória de nitrato disponível no solo (BODDEY *et al.*, 1986) e a fixação biológica do N₂ (INIGUEZ *et al.*, 2004). Entre esses mecanismos, o aumento do sistema radicular, estimulado pela presença de bactérias, através da produção de substâncias promotoras do crescimento radicular, pode resultar em maior absorção de minerais e de água (OKON; LABANDERA-GONZALEZ, 1994).

As características benéficas destas bactérias podem ser resumidas em: capacidade de penetrar na raiz das plantas, antagonismo a agentes patogênicos, associação com várias gramíneas e com não gramíneas (morango, tabaco, café e outras), produção de hormônios promotores de crescimento e desenvolvimento, baixa sensibilidade às variações de temperatura e ocorrência em todos os tipos de solo e clima (ARAÚJO, 2008).

É importante salientar que o processo de fixação biológica por essas bactérias em associação com gramíneas supre apenas parcialmente as necessidades das plantas em nitrogênio (HUNGRIA *et al.*, 2011). Não é recomendado a substituição total da adubação nitrogenada por bactérias diazotróficas não simbióticas.

2.6. Molibdênio

O molibdênio (Mo) tem sua principal função associada ao metabolismo do nitrogênio (N), e relaciona-se às enzimas redutase do nitrato e nitrogenase, de modo que os sintomas de deficiência se confundem com aqueles do nitrogênio (MARSCHNER, 1995). É capaz de mediar diversas reações de oxirredução nos sistemas biológicos (SRIVASTAVA, 1997). Participa como co-fator de enzimas redutase do nitrato, a oxidase da xantina, a oxidase de aldeído e a oxidase de sulfeto. A deficiência de Mo provoca redução na concentração de clorofilas nas folhas, acarretando decréscimo de fotossíntese e prejuízo no metabolismo do N, tendo como consequência o acúmulo de nitrato no tecido das plantas (BORKET, 1989). Portanto, qualquer deficiência desse elemento pode comprometer o metabolismo do N, diminuindo o rendimento das culturas.

O N absorvido pelas plantas na forma de nitrato (NO_3^-) é reduzido a amônia (NH_3), possibilitando assim sua assimilação. A primeira reação do processo redutivo é catalisada pela redutase do nitrato, sendo o Mo um cofator dessa enzima que reduz o NO_3^- a nitrito (NO_2^-). O (NO_2^-) é reduzido a amônia e assimilado na forma orgânica por meio do sistema glutamina sintetase glutamina oxoglutarato unida transferase (GS-GOGAT) com síntese de aminoácidos e, posteriormente, de proteínas, clorofila e outros compostos (CRAWFORD *et al.*, 1989). Assim, a produção de metabólitos nitrogenados (aminoácidos e proteínas) é afetada pela deficiência de Mo o uma vez que ocorre decréscimo na atividade da redutase do nitrato na ausência do cofator. Com isso, ocorre diminuição na síntese de aminoácidos e, conseqüentemente, de proteínas. Dessa forma, o Mo exerce papel direto no crescimento e desenvolvimento das plantas (VALETINI *et al.*, 2005).

As respostas à adubação molibídica estão relacionadas ao requerimento de Mo por vários tipos de molibdoenzimas (Mo-enzimas) presentes nas plantas. Essas Mo-enzimas podem estar envolvidas na redução e assimilação do N (nitrato redutase, NR), fixação do N (nitrogenase), catabolismo de purinas (xanthine dehydrogenase/oxidase), síntese de ácido abscísico, ABA, e ácido indol-3 acético (aldeído oxidase, AO) e metabolismo do enxofre (sulfite oxidase, SO) (KAISER *et al.*, 2005). Destas as de maior relevância para as plantas estão a nitrato redutase e a nitrogenase (HAMLIN, 2007).

O Mo utilizado pelas plantas pode ser originado do próprio solo ou resultante da aplicação de produtos químicos e/ou orgânicos que o contenham em sua composição (PEREIRA *et al.*, 2012). Geralmente, as fontes de micronutrientes de produtos químicos variam de modo considerável na sua forma física, reatividade química, custo, teor do nutriente

e eficiência agronômica. O fornecimento do fertilizante molíbdico às plantas tem sido feito de três formas principais: aplicação direta no solo, aplicação foliar e aplicação direta na semente (PEREIRA, 2010).

O Mo interfere diretamente no crescimento e desenvolvimento do milho e conseqüentemente, na produção de grãos, por meio do metabolismo do N (PEREIRA *et al.*, 1999). A faixa crítica de concentração de Mo no milho é de 0,1 a 0,2 mg kg⁻¹ (DIOS; BROYER, 1965). No milho, a deficiência de Mo encurta os internós, reduz a área foliar e causa o desenvolvimento de clorose nas folhas (AGARWALA *et al.*, 1978). O manejo adequado da adubação com molibdênio é imprescindível para o melhor aproveitamento da adubação nitrogenada (FERREIRA *et al.*, 2001).

REFERÊNCIAS BIBLIOGRÁFICAS

AGARWALA, S. C.; SHARMA, C. P.; FAROOQ, S.; CHATTERJEE, C. Effect of molybdenum deficiency on the growth and metabolism of corn plants raised in sand culture. **Canadian Journal of Botany**, v. 56, n. 16, p. 1905–1909, 1978.

ALVES, B. J. R.; ZOTARELLI, L., FERNANDES, F. M.; HECKLER, J. C.; MACEDO, R. A. T.; BODDEY, R. M.; JANTALIA, C. P.; URQUIAGA, S. Fixação biológica de nitrogênio e fertilizantes nitrogenados no balanço de nitrogênio em soja, milho e algodão (EMBRAPA AGROBIOLOGIA). **Pesquisa Agropecuária Brasileira**, v.41, n.3, p.449-456, 2006.

ALVES, G. C. **Efeito da Inoculação de Bactérias dos Gêneros *Herbaspirillum* e *Burkholderiana* Cultura do Milho**. 2007. 53 p. (Dissertação) - Programa de Pós-Graduação em Agronomia, Universidade Federal Rural do Rio de Janeiro, Seropédica, 2007. Disponível em: < <https://tede.ufrrj.br/jspui/handle/tede/267>>. Acesso em 30 de setembro de 2020

ANDRADE, A. C.; FONSECA, D. M.; QUEIROZ, D. S.; SALGADO, L. T.; CECON, P. R. Adubação nitrogenada e potássica em capim-elefante (*Pennisetum purpureum schum.* cv. napier). **Ciência e Agrotecnologia**, p.1643-1651, 2003.

ARAÚJO, S.C; **Realidade e perspectivas para o uso de *Azospirillum* na cultura do milho**. Piracicaba: IPNI – International Plant Nutrition Institute Brazil. 32p. (IPNI. Informações Agronômicas, 122). 2008.

BALDANI, V.L.D.; OLIVEIRA, E.; BOLOTA, E.; BALDANI, J.L.; KIRCHHOF, G.; DÖBEREINER, J. *Burkholderia brasiliensis* ssp. nov. uma nova espécie de bactéria diazotrófica endofítica. **Anais da Academia Brasileira de Ciência**. v. 69, p. 116. 1997.

BARTCHECHEN, A.; FIORI, C. C. L.; WATANABE, S. H.; GUARIDO, R. C. Efeito da inoculação de *Azospirillum brasiliense* na produtividade da cultura do milho (*Zea mays* L.). **Campo Digital**. v.5, n. 1, p.56-9, 2010.

BASHAN, Y.; HOLGUIN, G; DE-BASHAN, L.E. *Azospirillum* - plant relations physiological, molecular, agricultural, and environmental advances (1997-2003). **Canadian Journal of Microbiology**, v.50, n. 8, p.521-577, 2004.

BASHAN, Y.; HOGUIN, G. *Azospirillum* - plant relationship: Environmental and physiological advances (1990-1996). **Canadian Journal of Microbiology**. v.43, n. 2, p.103-121, 1997.

BASI, S. **Associação de *Azospirillum brasilense* e de nitrogênio em cobertura na cultura do milho**. 2013, 50 p. (Dissertação) – Programa de Pós-Graduação em Agronomia, Universidade Federal do Centro-Oeste, Guarapuava, 2013.

BELARMINO, M. C. J.; PINTO, J. C.; ROCHA, G. P.; FERREIRA NETO, A. E.; MORAIS, A. R. de. Altura de perfilho e rendimento de matéria seca de capim-tanzânia em função de diferentes doses de superfosfato simples e sulfato de amônio. **Ciência e Agrotecnologia**, v. 27, n. 4, p. 879-885, 2003.

BODDEY, R. M., BALDANI, V. L. D., BALDANI, J. I., DÖBEREINER, J. Effect of inoculation of *Azospirillum* spp. on the nitrogen assimilation of field grown wheat. **Plant and Soil**, v. 95, n. 1, p.109-121. 1986.

BOTREL, M. A.; ALVIM, M. J.; MARTINS, C. E. Aplicação de nitrogênio em acessos de *Brachiaria*. 2. Efeito sobre os teores de proteína bruta e minerais. **Pasturas Tropicais**, v. 12, n. 2, p. 7-10, 1999.

BORKET, C. M. Micronutrientes na planta. In: BILL, L. T. e ROSOLEM, C. A. (Ed.) **Interpretação de análise química de solo e planta para fins de adubação**. Ed. Fundação de Estudos e Pesquisas Agrícolas e Florestais, p. 309-329, 1989.

BREDA, F. A. D. F., ALVES, G. C., LOPEZ, B. D. O., ARAGÃO, A. R., ARAUJO, A. P., REIS, V. M. Inoculation of diazotrophic bacteria modifies the growth rate and grain yield of maize at different levels of nitrogen supply. **Archives of Agronomy and Soil Science**. v. 66, n. 14, p. 1948–1962. 2020.

BREDEMEIER, C.; MUNDSTOCK, C. M. Regulação da absorção e assimilação do nitrogênio nas plantas. **Ciência Rural**, v. 30, n. 2, p. 365-372, 2000.

BURRIS, R.H. Advances in biological nitrogen fixation. **Journal of Industrial Microbiology & Biotechnology**, v. 22, p.381-393, 1999.

CARVALHO, R. P.; VON PINHO, R. G.; DAVIDE, L. M. C. Eficiência e uso do nitrogênio em híbridos experimentais de milho do programa de melhoramento da Universidade Federal do Tocantins. **Revista Brasileira de Milho e Sorgo**, v. 10, n. 2, p. 108-120, 2011.

CONAB. COMPANHIA NACIONAL DE ABASTECIMENTO. **Acompanhamento da safra brasileira: grãos. 12º Levantamento**. Brasília, DF: 2020. Disponível em: <[https://www.conab.gov.br/info-agro/safras/graos/boletim-da-safra-de-
graos/boletim_graos_setembro_2020.pdf](https://www.conab.gov.br/info-agro/safras/graos/boletim-da-safra-de-graos/boletim_graos_setembro_2020.pdf)> Acesso em 31 de maio de 2021.

CORREA, O. S.; ROMERO, A. M.; SORIA, M. A.; DEESTRADA, M. *Azospirillum brasilense* – plant genotype interactions modify tomato response to bacterial diseases, and root and foliar microbial communities. In: CASSÁN, F.D.; GARCIA DE SALAMONE, I. (Ed.) **Azospirillum sp.: cell physiology, plant interactions and agronomic research in Argentina**. Argentina: Asociación Argentina de Microbiología, p.87-95, 2008.

CRAWFORD, N. M.; CAMPBELL, W. H.; DAVIS, R. W. Nitrate reductase from squash: DNA cloning and nitrate regulation. **Proceedings of the Nacional Academy of Sciences**, v.83, n. 21, p.8073-8076, 1986.

DARTORA, J.; GUIMARÃES, V. F.; MARINI, D.; SANDER, G. Adubação nitrogenada associada à inoculação com *Azospirillum brasilense* e *Herbaspirillum seropedicae* na cultura do milho. **Revista Brasileira de Engenharia Agrícola e Ambiental**.v.17, p. 1023-1029, 2013.

DIOS, R.V.; BROYER, T.C. Deficiency symptoms and essentiality of molybdenum in corn hybrids. **Agrochimica**, v. 9, n. 3, p. 273, 1965.

DOBBELAERE, S.; VANDERLEYDEN, J.; OKON, Y. Plant growth-promoting effects of diazotrophs in the rhizosphere. **Critical Reviews in Plant Science**, v. 22, n. 2, p.107-149, 2003.

DÖBEREINER, J. History and new perspective of diazotrophs in association with non-leguminous plants. **Symbiosis - rehovot** subscription magazine v.13, n.1, p.1-13, 1992.

ECKERT, B., WEBER, O. B., KIRCHHOF, G., HALBRITTER, A., STOFFELS, M. HARTMANN, A. *Azospirillum doebereinae* sp. nov. a nitrogen fixing bacterium associated with the C4-grass Miscanthus. **International Journal of Systematic and Evolutionary Microbiology**, v. 51, n.1, p.17–26, 2001.

FAGAN, E. B.; MEDEIROS, S. L. P.; MANFRON, P. A.; CASAROLI, D.; SIMON, J.; DOURADO, D. N.; MÜLLER, L. Fisiologia da fixação biológica do nitrogênio em soja - Revisão. **Revista da FZVA**, v. 14, n. 1, p. 89-106, 2007.

FAGERIA, N. K. Otimização da eficiência nutricional na produção das culturas. **Revista Brasileira de Engenharia Ambiental**, v.2, n.1, p 6-16, 1998.

FAGERIA, N. K.; SLATON, N. A.; BALIGAR, V. C. Nutrient management for improving lowland rice productivity and sustainability. **Advances in Agronomy**, v. 80, p. 63-152, 2003.

FAGERIA, N. K.; SANTOS, A. B.; CUTRIM, V. A. Produtividade de arroz irrigado e eficiência de uso do nitrogênio influenciadas pela fertilização nitrogenada. **Pesquisa Agropecuária Brasileira**, v. 42, n. 7, p. 1029-1034, 2007.

FARINELLI, R.; LEMOS, L. B. Produtividade e eficiência agrônômica do milho em função da adubação nitrogenada e manejos do solo. **Revista Brasileira de Milho e Sorgo**, v. 9, n. 2, p. 135-146, 2010.

FERNANDES, F. C. S.; BUZETTI, S.; ARF, O.; ANDRADE, A.C. Doses, eficiência e uso de nitrogênio por seis cultivares de milho. **Revista Brasileira de Milho e Sorgo**, v.4, n. 2, p. 195-204, 2005.

FERNANDES, F. C. S.; LIBARDI, P. L. Percentagem de recuperação de nitrogênio pelo milho, para diferentes doses e parcelamentos do fertilizante nitrogenado. **Revista Brasileira de Milho e Sorgo**, v.6, n. 3, p. 285-296, 2007.

FERNANDES, F. C. S.; LIBARDI, P. L.; TREVELIN, P. C. O. Parcelamento da adubação nitrogenada na cultura do milho e utilização do N residual pela sucessão aveia preta-milho. **Ciência Rural, Santa Maria**, v. 38, n. 4, p. 1138-1141, 2008.

FERNANDÉZ, J. E.; MURILLO, J. M.; MORENO, F.; CABRERA, F.; FERNANDÉZ-BOY, E. Reducing fertilization for maize in southwest Spain. **Communications in Soil Science and Plant Analysis**, v. 29, n. 19-20, p. 2829-2840, 1998.

FERREIRA, A. C. D. B., ARAÚJO, G. A. D. A., PEREIRA, P. R. G.; CARDOSO, A. A. Corn crop characteristics under nitrogen, molybdenum and zinc fertilization. **Scientia Agricola**, v. 58, n. 1, p. 131-138, 2001.

FIDELIS, R. R.; MIRANDA, G. V.; SANTOS, I. C.; GALVÃO, J. C. C.; PELUZIO, J. M.; LIMA, S. O. Fontes de germoplasma de milho para estresse de baixo nitrogênio. **Pesquisa Agropecuária Tropical**, v. 37, n. 3, p. 147-153, 2007.

FIGUEIREDO, M. V. B.; JUNIOR, M. A. L.; MESSIAS, A. S.; MENEZES, R. S. C. **Potential Impact of biological nitrogen fixation and organic fertilization on corn growth and yield in low external input systems**. In: DANFORTH, A.T. (Ed.). *Corn crop production growth, fertilization and yield*. New York: Nova Science Publisher, p.227-255, 2009.

FORDE, B. G. Nitrogen signalling pathways shaping root system architecture: an update. **Current Opinion in Plant Biology**. v 21, p. 30–36, 2014.

GAJU, O.; ALLARD, V.; MARTRE, P.; LE GOUIS, J.; MOREAU, D.; BOGARD, M.; HUBBART, S; FOULKES, M. J. Nitrogen partitioning and remobilization in relation to leaf senescence, grain yield and grain nitrogen concentration in wheat cultivars. **Field Crops Research**, v. 155, p.213-223, 2014. DOI: <http://dx.doi.org/10.1016/j.fcr.2013.09.003>.

GANAPATHY, B. A.; SAVALGI, V. P. Effect of micronutrients on the performance of *Azospirillum brasilense* on the nutrient uptake, growth and yield in maize crop. **Karnataka Journal of Agricultural Sciences**, v.19, n. 1, p.66-70, 2006.

HAMLIN, R. L. **Molybdenum**. In: BARKER, A. V.; PILBEAM, D. J. Handbook of plant nutrition. London: Taylor e Francis, 613 p. 2007.

HIREL, B.; LE GOUIS, J.; NEY, B.; GALLAIS, A. The challenge of improving nitrogen use efficiency in crop plants: towards a more central role for genetic variability and quantitative genetics within integrated approaches. **Journal Experimental Botanic**. v. 58, n. 9, p. 2369-2387, 2007.

HUNGRIA, M. **Inoculação com *Azospirillum brasilense*: inovação em rendimento a baixo custo**. Londrina: Embrapa Soja, (EMBRAPA SOJA. Documentos, 325), 37p. 2011.

INIGUEZ, A. L.; DONG, Y.; TRIPLETT, E. W. Nitrogen fixation in wheat provided by *Klebsiella pneumoniae* 342. Molec. **Plant Microbiology**, v.17, n. 10, p.1078-1085. 2004.

KAISER, B. et al. The Role of Molybdenum in Agricultural Plant Production. **Annals of Botany**, v. 96, n. 5, p. 745-754, 2005.

KOLCHINSKI, E. M; SCHUCH, L. O. B. Eficiência no uso do nitrogênio por cultivares de aveia branca de acordo com a adubação nitrogenada. **Revista Brasileira de Ciência do Solo**, v. 27, p. 1033-1038, 2003.

KONG, L.; XIE, Y.; HU, L.; FENG, B.; LI, S. Remobilization of vegetative nitrogen to developing grain in wheat (*Triticum aestivum* L.). **Field Crop Research**. v.196, p.134-144, 2016.

LE GOUIS, J., BEGHIN, D., HEUMEZ, E.; PLUCHARD, P. Genetic differences for nitrogen uptake and nitrogen utilization efficiencies in winter wheat. **European Journal of Agronomy**, v. 12, n. 3-4, p. 163-173, 2000.

MAGALHÃES, P. C.; DURÃES, F. O. M.; CARNEIRO, N. P.; PAIVA, E. **Fisiologia da planta de milho**. Embrapa Milho e Sorgo-Circular Técnica (INFOTECA-E), 2006.

MARSCHNER, H. **Mineral nutrition of higher plants**. New York, Academic Press, 889 p. 1995.

MENDES, E. G. R. **Uso do inibidor de urease para aumentar a eficiência do nitrogênio na cultura do milho**. 2016, 46 p. (Dissertação) - Programa de Pós-Graduação em Agroecologia, Universidade Estadual do Maranhão, São Luís. 2016. Disponível em: <<http://repositorio.uema.br/123456789/199>>. Acesso em: 24 de setembro de 2019.

MENGEL, K.; KIRKBY, E.A. **Principles of plant nutrition**. Bern: International Potash Institute, p. 687-695, 1987.

MOREIRA, F. M. S.; SILVA, K.; NÓBREGA, R. S. A.; CARVALHO, F. Bactérias diazotróficas associativas: diversidade, ecologia e potencial de aplicações. **Comunicata Scientiae**, v. 1, n. 2, p. 74-99, 2010.

MOREIRA, J. C. **Acúmulo de matéria seca e de nutrientes na cultura do milho verde**. 2015. 56 p. (Dissertação) - Pós-Graduação em Fitotecnia, Universidade Federal Rural do Semi-Árido, 2015. Disponível em: <<http://repositorio.ufersa.edu.br/handle/tede/106>>. Acesso em: 15 de janeiro de 2020.

MÔRO, G. V.; FRITSCHÉ, N. R. **Importância e uso do milho no Brasil**. In: BORÉM, A.; GALVÃO, J. C. C.; PIMENTEL, M. A. (Eds.). *Milho: do plantio à colheita*. 1º ed. Viçosa: UFV: p. 09-25, 2015.

MOLL, R. H.; KAMPRATH, E. J.; JACKSON, W. A. Analysis and interpretation of factors which contribute of efficiency of nitrogen utilization. **Agronomy Journal**, v.74, n. 3, p. 562-564, 1982.

MYLONA, P; PAWLOWSKI, K.; BISSELING T. Symbiotic Nitrogen Fixation. **The Plant Cell**, v.7, n. 7, p.869-885, 1995.

NOVAKOWISKI, J. H.; DANDINI, I. E.; FALBO, M. K.; MORAES, A.; CHENG, N. C. Efeito Residual da Adubação Nitrogenada e Inoculação de *Azospirillum brasilense* na Cultura do Milho. **Semina: Ciências Agrárias**, v. 32, n.1, p. 1687-1698, 2011.

OKON, Y.; LABANDERA-GONZALES, C. A. Agronomic applications of *Azospirillum*: an evaluation of 20 years worldwide field inoculation. **Soil Biology and Biochemistry**, v.26, n. 12, p.1591-1601. 1994.

OKUMURA, R. S.; MARIANO, D. C.; ZACCHEO, P. V. C. Uso de fertilizante nitrogenado na cultura do milho: uma revisão. **Pesquisa Aplicada & Agrotecnologia**, v. 4, n. 2, p. 226–244, 2011.

PEREIRA, F. R. da S. **Doses e formas de aplicação de molibdênio na cultura do milho**. 2010. 141 p. (Tese) Programa de Pós-Graduação em Agronomia, Universidade Estadual Paulista, Faculdade de Ciências Agrônomicas, 2010. Disponível em: <<http://hdl.handle.net/11449/99959>>. Acesso em: 14 de março de 2020.

PEREIRA, S. L., ARAÚJO, G. D. A., SEDIYAMA, C. S., VIEIRA, C.; MOSQUIM, P. R. Efeitos da adubação nitrogenada e molíbdica sobre a cultura do milho. **Ciência e Agrotecnologia**, v. 23, n. 4, p.:4 - 44, 1999.

PEREIRA, F. R. da. S.; BRACHTVOGEL, E. L.; CRUZ, S. C. S, SILVIO JOSÉ BICUDO, S. J.; MACHADO, C. G.; PEREIRA, J. C. Qualidade fisiológica de sementes de milho tratadas com molibdênio. **Revista Brasileira de Sementes**, v. 34, n. 3 p. 450 - 456, 2012.

PERSELLO-CARINEAUX, F.; NUSSAUME, L.; ROBAGIA, C. Tales from the underground: molecular plant rhizobacteria interactions. **Plant Cell and Environment**, v. 26, n. 2, p. 189-99, 2003.

PICAZEVICZ, A. A. C.; KUSDRA, J. F.; MORENO, A. de L. Maize growth in response to *Azospirillum brasilense*, *Rhizobium tropici*, molybdenum and nitrogen. **Revista Brasileira de Engenharia Agrícola e Ambiental**, v. 21, n. 9, p.623-627, 2017.

SANGOI, L.; DA SILVA, L. M. M.; MOTA, M. R.; SCHMITT, F. P. A.; SOUZA, N. M.; GIORDANI, W.; SCHENATTO, D. E. Desempenho agrônômico do milho em razão do tratamento de sementes com *Azospirillum* sp. e da aplicação de doses de nitrogênio mineral. **Revista Brasileira de Ciência do Solo**, v. 39, p.1141-1150, 2015.

RITCHIE, S.W.; HANWAY, J.J.; BENSON, G.O. Como a planta de milho se desenvolve. POTAFOS. **Informações Agronômicas**, n.103, p.1-19, 2003.

SÁ JÚNIOR, A.; **Comportamento agrônômico do milho em resposta ao modo de aplicação e concentrações de *Azospirillum brasilense***. 2012. 51 p. (Dissertação) – Programa de Pós-Graduação em Agronomia, Instituto de Ciências Agrárias, Universidade Federal de Uberlândia, Uberlândia, 2012. Disponível em: <<https://repositorio.ufu.br/handle/123456789/12182>>. Acesso em: 07 de março de 2020.

SCIVITTARO, W. B.; MURAOKA, T.; BOARETTO, A. E.; TRIVELIN, P. C. O. Utilização de nitrogênio de adubos verdes e mineral pelo milho. **Revista Brasileira de Ciência do Solo**, v.24, n.4, p.917-926, 2000.

SILVA, G. F.; OLIVEIRA, F. H. T. de.; PEREIRA, R. G.; SILVA, P. S. L. DIÓGENES, T. B. A.; SILVA, A. R. da C. Doses de nitrogênio e fósforo para produção econômica de milho na Chapada do Apodi, RN. **Revista Brasileira de Engenharia Agrícola e Ambiental**. v. 18, n. 12, p. 1247-1254, 2014.

SILVA, T. F.; MELLONI, R. Densidade e diversidade fenotípica de bactérias diazotróficas não simbióticas em solos da Reserva Biológica Serra dos Toledos, Itajubá (MG). **Revista Brasileira de Ciência do Solo**, v. 35, n. 2, p. 359-371, 2011.

SOUZA, C. F. D. **Desempenho agrônômico e eficiência de utilização de nitrogênio por cultivares de milho**. 2017. 51 p. (Tese) - Programa de Pós-Graduação em Fitotecnia da Universidade Federal Rural do Semiárido. Disponível em: <<http://bdtd.ufersa.edu.br/handle/tede/728>> Acesso em: 15 de outubro de 2017.

SOUZA, S. R.; FERNANDES, M. S. In: FERNANDES, M. S. **Nutrição Mineral de Plantas**. Sociedade Brasileira de Ciência do Solo. p. 215-252, 2006.

SRIVASTAVA, P. C. Biochemical significance of molybdenum in crop plants. In: GUPTA, U. C. **Molybdenum in agriculture**. New York: Cambridge University Press, p. 47-70, 1997.

STEENHOUDT, O.; VANDERLEYDEN, J. *Azospirillum*, a free-living nitrogen-fixing bacterium closely associated with grasses: genetic, biochemical and ecological aspects. **FEMS Microbiology Reviews**, v.24, n. 4, p.487-506, 2000.

TAIZ, L.; ZEIGER, E. **Fisiologia Vegetal**. Trad. MASTROBERTI, A. A. et. al. 6° ed.; Porto Alegre: Artmed, 2017, 888p.

USDA. **Grain: World Markets and Trade**. United States Department of Agriculture. Foreign Agricultural Service. p. 1-56. 2017. Acesso em 28 de setembro de 2017. Disponível em: <<https://apps.fas.usda.gov/psdonline/app/index.html#/app/downloads>>. Acesso em: 20 de março de 2020.

VALENTINI, L.; CUNHA, C. F.; DOS SANTOS, M. F. Teor de nitrogênio foliar e produtividade de três cultivares de milho (*Zea mays* L.) submetidos às adubações nitrogenada e molíbdica. **Revista Ceres**, v.52, p.567-577, 2005.

WANG, X.; XING, Y. Effects of irrigation and nitrogen on maize growth and yield components. **Global Changes and Natural Disaster Management: Geo-information Technologies**. v. 1, p.63-74, 2017. DOI:

WEBER, O. B.; BALDANI, J. I.; DÖBEREINER, J. Bactérias diazotróficas em mudas de bananeira. **Pesquisa Agropecuária Brasileira**, v. 35, n.11, p. 2277- 2285, 2000.

WILLIAMS, L. E.; MILLER, A. J. Transporters responsables for the uptake and Partitioning of nitrogenous solutes. **Annual Review Plant Physiology and Molecular Biology**, v. 52, n. 1, p. 659-688, 2001.

YAMADA, T.; ABDALLA, S.R.S. Como melhorar a eficiência da adubação nitrogenada do milho. **Informações Agronômicas**, v.91, p.1-5, 2000.

ZAIED, K. A.; EL-HADY, A. H.; AFIFY, A. H.; NASSEF, M. A. Yield and nitrogen assimilation of winter wheat inoculated with new recombinant inoculants of rhizobacteria. **Journal of Biological Sciences**, v.4, p.344-358. 2003.

The effects of nitrogen, *Azospirillum brasilense* and molybdenum on the grain yield and nitrogen use efficiency on maize plants in the sub-humid tropics of Brazil.

CAPÍTULO II

1 **The effects of nitrogen, *Azospirillum brasilense* and molybdenum on**
2 **the grain yield and nitrogen use efficiency on maize plants in the sub-**
3 **humid tropics of Brazil.**

4 **Marcelo Marinho Viana^a, * and Heder Braun^a**

5 ^a*Postgraduate Program in Agroecology, Department of Plant Science and Plant Disease, Maranhão*

6 *State University, São Luís, Maranhão, Brazil.*

7 *** Corresponding author:**

8 *E-mail address: marceloviana.91@gmail.com (Marcelo Marinho Viana)*

9 **Highlights**

- 10 • The combination (*A. brasilense*, Mo and N) increased 5.1% nitrogen use
11 efficiency;
- 12 • Application methods *A. brasilense* on seed increased 13.4% nitrogen use
13 efficiency;
- 14 • Molybdenum and nitrogen increased 17.8% grain yield;

15

16

17

18

19

20

21

22

23

24 ABSTRACT

25 The objective this study was to evaluate the effectiveness of inoculation methods of
26 *A. brasilense*, N rate and molybdenum on yield and N use efficiency in maize plants
27 cultivated in the sub-humid tropical regions of Brazil. Three field trials were conducted
28 in a randomized complete block design with four replicates and eight treatments. The
29 treatments were 140 kg ha⁻¹ of N, 90 g ha⁻¹ of Mo and inoculation methods of *A.*
30 *brasilense* (seed and leaf). In experiment 2, all treatments did affect grain yield, 100 -
31 seed weight, harvest index, leaf chlorophyll index (LCI), stem diameter, insertion height
32 of the first ear (IHFE) and shoot dry matter. In experiment 3, all treatments did affect
33 grain yield, nitrogen use efficiency (NUE), harvest index, leaf chlorophyll index (LCI),
34 stem diameter, insertion height of the first ear (IHFE), plant height and shoot dry matter.
35 The inoculation with *A. brasilense*, N and Mo showed the potential to be used, and the
36 results showed that Mo is the limiting factor for grain yield and NUE.

37

38 **Keywords:** nitrogen fertilization, biological nitrogen fixation, plant growth-promoting
39 rhizobacteria, *Zea mays*.

40 1. INTRODUCTION

41 Nitrogen (N) is the most significant agricultural input for crops to achieve high
42 yield, is consumed by the crop roots throughout the growing season (Wang and Xing,
43 2017; Picazevicz, Krudra and Moreno, 2017; Galindo et al. 2017; Bloch et al. 2020), but
44 excessive N inputs make it a difficult problem for optimal use of N (Norr, 2017). Excess
45 or low supply of the nutrients will result in reduced nitrogen use efficiency (NUE) and
46 cause significant losses in grain yield and grain quality (Haroon et al, 2019).

47 Nitrogen is considered a major limiting factor for maize (*Zea mays* L.) grain yield
48 because it is an essential component of all proteins and enzymes, nucleic acids that make
49 up DNA, and chlorophyll that enables the process of photosynthesis in plants (Leghari et

50 al. 2016). Low yields are generally attributed to low fertility soils that require high
51 fertilizer inputs for optimal productivity (Martins et al, 2018). In this context cultivation
52 of maize in tropical soil from the Amazonian periphery do not adequately supply the
53 plant's demand for nitrogen.

54 Agronomic management for improved nitrogen use efficiency (NUE), the use of
55 agroecological practices is the key to increase maize production while reducing
56 environmental pollution. One possibility such as inoculation by plant growth-promoting
57 bacteria (PGPB) can represent a sustainable alternative for increase nutrient use
58 efficiency in tropical agriculture (Perreira et al. 2020). Inoculation technology with PGPB
59 has been presented worldwide as an important tool for reaching sustainability in
60 agriculture due to its low environmental and production costs compared with industrial
61 inputs (Oliveira et al. 2017).

62 PGPB colonize rhizosphere or plant root and improve plant health and growth.
63 Some of the most important plant growth promoting bacteria activity include biological
64 nitrogen fixation (BNF)(Pankiewicz et al., 2019), production of indolic compounds and
65 siderophores, increase on 1-aminocyclopropane-1-carboxylate deaminase activit
66 (Ambrosini and Passaglia, 2017), solubilization of mineral phosphates (Ludueña et al.,
67 2018; Qi et al., 2018) and production of phytohormones, such as salicylic acid,
68 gibberellins, cytokinins and indole-3-acetic acid (IAA) (Cassán and Diaz-Zorita, 2016;
69 Fukami et al., 2017; Dahal et al., 2017; Gouda et al., 2018). The diazotroph *Azospirillum*
70 *brasilense* (free-living diazotrophic bacteria) is considered a model PGPB, and a great
71 amount of information regarding the physiology of its growth and development has been
72 published (Fendrik et al., 1995; Cassán et al., 2015, Cassán et al, 2020). Several positive
73 results in the development and productivity of corn have been reported with inoculation

74 with *Azospirillum brasilense* (strains Ab-V5 and Ab-V6) in tropical conditions (Martins
75 et al., 2018; Oliveira et al., 2018; Galindo et al., 2019).

76 Molybdenum is the micronutrient required in the least amount by plants.
77 However, there is a close relationship between the supply of Mo, the activity of nitrate
78 reductase, and the growth of plants. (Kirkby and Römheld, 2004). Therefore, the supply
79 of Mo is closely associated with the utilization and metabolism of N. Nitrogen utilization
80 by maize can be potentiated by molybdenum (Mo), since it is a constituent of enzymes
81 that allow the assimilation of this macronutrient via biological fixation by diazotrophic
82 microorganisms (nitrogenase) and/or N fertilization (nitrate reductase) (Picazevicz et al,
83 2017). The increase in maize growth and production has already been observed with the
84 use of Mo combined with *Azospirillum brasilense* (Ganapathy and Savalgi, 2006) and N
85 fertilizer (Valentini et al., 2005)

86 The combined use of chemical and biological inputs in the cultivation of non-
87 legumes can contribute to reduce costs and optimize production (Picazevicz et al, 2017).
88 Less is known about the effects of inoculation methods of *A. brasiliense*, N rates and
89 molybdenum on physiological characteristics, yield, and N use efficiency on maize plants
90 grown in the sub-humid tropical regions of Brazil. Thus, we evaluated the effectiveness
91 of inoculation methods of *A. brasiliense*, nitrogen and molybdenum on yield and N use
92 efficiency in maize plants cultivated in the sub-humid tropical regions of Brazil.


93 **2. MATERIALS E METHODS**

94

95 *2.1. Field site description*

96 Three field trials were conducted in the municipality of São Luís (2°30'S,
97 44°18'W, 24 m above sea level), State of Maranhão, Brazil. The first trial was established
98 from February 2018 to May 2018, the second trial from October 2018 to January 2019

99 and third trial from March 2019 to July 2019. The region has a hot, semi-humid,
 100 equatorial climate, with mean annual rainfall of 2,200 mm and two well-defined seasons:
 101 a rainy season from January to June, and a dry season with pronounced water deficits
 102 from July to December. The climatic conditions during the trials are shown in Figure 1.
 103



104

105 **Figure 1.** Rainfall (mm) and maximum and minimum temperatures obtained from the
 106 data base National Institute of Meteorology of Brazil (INMET) during the corn cultivation
 107 (all seasons) in the period from January 2018 to July 2019.

108

109 The trials were conducted under a no-tillage system. The field had not been
 110 cultivated with any agricultural crop for at least 10 years (2003-2013). Maize and cowpea
 111 (*Vigna unguiculata* L. Walp) had been cultivated in the field after 2013. The remaining
 112 straws from previous maize and cowpea crops were left on the soil surface. The soil in
 113 this area is a Typic Hapludult (Soil Survey Staff, 1999), with pH in CaCl₂ (soil:solution

114 ratio of 1:2.5) = 5.3; organic matter = 5 g dm⁻³; P (resin) = 31 mg dm⁻³; K⁺ = 1.7 mmolc
 115 dm⁻³; Ca²⁺ = 24 mmolc dm⁻³; Mg²⁺ = 15 mmolc dm⁻³; H⁺ + Al³⁺ = 19 mmolc dm⁻³; sum
 116 of bases = 40.7 mmolc dm⁻³; cation exchange capacity at pH 7 = 59.7 mmolc dm⁻³; base
 117 saturation = 68%; and coarse sand = 260 g kg⁻¹, fine sand = 640 g kg⁻¹, silt = 20 g kg⁻¹,
 118 and clay = 120 g kg⁻¹ and texture sandy loam.

119 No mechanical soil preparation was carried out before the maize sowing. At 30
 120 days before sowing in 2017 was applied 1.0 Mg ha⁻¹ of dolomitic limestone (32% CaO,
 121 15% MgO, and total neutralizing power of 91%) without incorporation.

122 2.1. Experimental design, treatment, and field management

123 The experimental design was a randomized complete block design with four
 124 replicates and eight treatment (Table 1). The treatments were composed from one N rate
 125 of 140 kg ha⁻¹, one Mo rate of 90 g ha⁻¹ and inoculation methods of *A. brasilense* (seed
 126 and leaf). Each plot consisted of four 4 m-long rows, spaced 0.8 m apart. The outer rows
 127 of each plot and 0.5 m from each end of the rows were used as borders.

128 **Table 1.** Treatment arrangement of all seasons 2018/2019 with *Azospirillum brasilense*
 129 (seeds and leaf), nitrogen and molybdenum rates.

Treatments	Nitrogen (kg/ha ⁻¹)		Molybdenum (g/ha ⁻¹)	Azospirillum seeds (ml/ha ⁻¹)	Azospirillum leaf (ml/ha ⁻¹)
	Sowing	Topdressing	Topdressing	Sowing	Topdressing
1	-	-	-	-	-
2	40	100	-	-	-
3	40	100	-	100	-
4	40	100	-	-	200
5	40	100	90	-	-
6	40	100	90	100	-
7	40	100	90	-	200
8	40	100	90	100	200

130

131 At sowing furrow, 40 kg ha⁻¹ of N (granulated urea), 80 kg ha⁻¹ of P₂O₅ (simple
 132 superphosphate), and 100 kg ha⁻¹ of K₂O (potassium chloride) were applied in all trials.
 133 Seeds of the maize simple hybrid PIONNER 30F35[®] was sown at a density of four seeds

134 per meter (5 plants m⁻²). Seedling emergence occurred between four and six days after
135 sowing for all trials. All plots were irrigated after the sowing with 15 mm of water, using
136 a drip tape system and 2-day intervals, for a good and uniform seedling emergence.
137 Approximately every 3 days without rainfall after seedling emergence, the plots were
138 irrigated with 15 mm of water (2-hour irrigation). The plants were irrigated when there
139 was no or low rainfall in the previous week. When there was a need to irrigate, we use
140 the drip tape with flat emitters inside, 16mm diameter, thickness from 0.6mm, spacing
141 from 20cm.

142 Topdressed N (100 kg ha⁻¹ of N) (granulated urea, with 46% of N) was manually
143 applied when the maize plants had six leaves completely expanded (V6) and was evenly
144 applied along the furrows at 10 cm away from the plants. All plots were immediately
145 irrigated after N fertilizer with approximately 15 mm of water to minimize ammonia
146 volatilization. Topdressed Mo (90 g ha⁻¹) (ammonium molybdate) applied when the maize
147 plants had six leaves completely expanded (V6) by spraying the solution, using a sprayer
148 equipped with one cone nozzle (XR 11002; Teejet®, Wheaton, USA), with a flow rate of
149 200 L ha⁻¹ of water. A plastic sheet was used to protect adjacent plots from unwanted
150 spray drift.

151 The seeds and leaves of maize plants were inoculated with *Azospirillum*
152 *brasilense*, using the commercial strains Ab-V5 and Ab-V6 (Nitro 1000 Gramíneas; Nitro
153 1000, Cascavel, Brazil) at rates of 100 mL and 200 mL of the liquid inoculant per hectare
154 (2×10⁸ CFU [colony forming unity] mL⁻¹), respectively. The seeds were microbiolized
155 one hour before sowing the crop, and the leaves were inoculated at V6 stage maize by
156 spraying the solution, using a sprayer equipped with one cone nozzle (XR 11002;
157 Teejet®, Wheaton, USA), with a flow rate of 200 L ha⁻¹ of water. A plastic sheet was
158 used to protect adjacent plots from unwanted spray drift.

159 Deltamethrin (5g ha⁻¹ active ingredient) was applied to control fall armyworm
160 (*Spodoptera frugiperda*), when needed. Weeds were controlled by manual hoeing until
161 the N topdressing application.

162 2.3. Measurements collected

163 The following nutritional evaluations were performed: leaf chlorophyll index
164 (LCI), measured indirectly using a portable non-destructive chlorophyll meter SPAD-502
165 (Minolta Co., Japan). The readings were performed in 5 plants per plot, and total N
166 concentration in leaves, collecting 2 leaves in the flowering stage were analyzed using the
167 Kjeldahl method (Tedesco et al., 1995).

168 The following productive component measurements were performed at 100 days
169 after emergence for all experiments, plant height, defined as the distance (cm) from the
170 ground level to the apex of the spike, at harvest time; stem diameter, insertion height of
171 the first ear (IHFE). Five plants from each plot were collected from two central rows to
172 determine plant dry weight and plant N concentration. These plants were separated into
173 grain and shoot plant, then oven-dried at 70 °C to constant weight and weighed. Grain
174 dry weight plus shoot dry weight represented the total plant dry weight. Total N
175 concentration of each fraction (grain and shoot plant) was analysed using the Kjeldahl
176 method (Tedesco et al., 1995) and the N content was calculated as the product of N
177 concentration by its dry weight. Grain N content plus shoot N content represents the plant
178 N content (aboveground). All ears from the central two rows of each plot were hand-
179 harvested. Grain yield was calculated and adjusted to 13% moisture content and 100-seed
180 weight. This moisture content is the minimum considered for the marketing of maize in
181 Brazil.

182 Based on the measurements of plant dry weight and N content, we calculated
183 harvest index dividing the grain dry weight by the plant dry weight and N harvest index

184 dividing the grain N content by the plant N content. The N use efficiency and its two
185 components were calculated according to Moll et al. (1982): i) N utilization efficiency
186 (kg kg^{-1} , kg of grain per kg of N extracted) = grain yield/plant N content, ii) N uptake
187 efficiency (kg kg^{-1} , kg of N in plant per kg of N applied) = plant N content/N rates, iii) N
188 use efficiency (kg kg^{-1} , kg of grain per kg of N applied) = $\text{NUpE} \times \text{NUE} = \text{grain yield/N}$
189 rates.

190 2.4. Statistical analyses

191 All data were initially tested for homogeneity of variance with O'Neill and
192 Mathews test and for normality using the Shapiro Wilk test. When the effect of treatment
193 was significant ($p < 0.05$) its sum of squares were partitioned into seven orthogonal
194 contrasts (below). Values were reported as means \pm standard deviation ($n=4$).

195 The following orthogonal contrasts are:

- 196 a) T1 vs. T2 to T8 (this contrast check the effect of N deficiency in maize plants)
- 197 b) T2 vs. T3 to T8 (this contrast check the effect of new production technology)
- 198 c) T3 and T4 vs T5 to T8 (this contrast check the effect of Azospirillum versus the
199 combination Mo and Azospirillum)
- 200 d) T3 vs. T4 (this contrast check the effect of Azospirillum on seed versus Azospirillum
201 on leaf, without Mo application)
- 202 e) T5 vs T6 to T8 (this contrast check the effect of Mo versus Azospirillum, regardless of
203 the form of application)
- 204 f) T6 vs T7 and T8 (this contrast check the effect of Azospirillum on seed versus
205 Azospirillum, regardless of the form of application)
- 206 g) T7 vs T8 (this contrast check the effect of Azospirillum on leaf versus combination
207 Azospirillum on seed with Mo application).

208 Statistical analyses were performed using the statistical Software R version 4.0.2 (R
209 Core Team, 2021) and the ExpDes.pt package (Ferreira; Calvacanti and Nogueira, 2014).

210 3. RESULTS

211 In experiment 1, treatments affected grain yield, 100-seed weight, nitrogen use
212 efficiency (NUE), leaf chlorophyll index (LCI), stem diameter, insertion height of the
213 first ear (IHFE) and shoot dry matter (Table 2). Treatments did not affect harvest index
214 ($p = 0.50$, mean = 63.56 ± 2.93 %) and plant height ($p = 0.24$, mean = 1.54 ± 0.11 m).

215 In experiment 2, treatments affected grain yield, 100 - seed weight, harvest index,
216 leaf chlorophyll index (LCI), stem diameter, insertion height of the first ear (IHFE) and
217 shoot dry matter, however, treatments did not affect NUE ($p = 0.06$, mean = 36.50 ± 3.67
218 kg kg^{-1}) and plant height ($p = 0.10$, mean = 1.64 ± 0.08 m).

219 In experiment 3, all treatments did affect grain yield, nitrogen use efficiency
220 (NUE), harvest index, leaf chlorophyll index (LCI), stem diameter, insertion height of the
221 first ear (IHFE), plant height and shoot dry matter, however, treatments did not affect 100
222 - seed weight ($p = 0.19$, mean = 22.49 ± 1.78 g).

223 Maize plants N deficiency was either 31% for grain yield, 4.2% LCI, 5.3 % stem
224 diameter, 5.9% IHFE, 4.4% shoot dry matter (Exp. 1), 23.6% grain yield, 14.7% LCI,
225 35.2% stem diameter, 13.8% IHFE, 4.8% shoot dry matter (Exp. 2), 39.3% grain yield,
226 17.5% harvest index, 28.9% LCI, 23.7% plant height, 10.5% stem diameter, 10.7% IHFE
227 (Exp. 3) greater compared when maize plants fertilized of nitrogen. However, 3.24%
228 shoot dry matter was increased with N deficiency in experiment 3 (1 vs T2 to T8, Table
229 3 and 4).

230 Maize plants fertilized with 140 kg ha^{-1} of nitrogen, 90 g ha^{-1} of Mo and inoculated
231 *A. brasilense* (seed and leaf) was either 3.9% stem diameter and 5.5% shoot dry matter

232 (Exp.1), 8.3% for 100 seed weight, 15.3% stem diameter (Exp.2) and 5.1% NUE and
233 2.9% IHFE (Exp. 3) greater compared when maize plants only fertilized with 140 kg ha⁻¹
234 of nitrogen. However, there was a reduction 6.3% for 100 - seed weight, 4.2% IHFE
235 (Exp. 1), 10.6% IHFE, 9.3% shoot dry matter (Exp. 2), 20.5% LCI, 4.8% stem diameter
236 and 8.5% shoot dry matter (Exp. 3) (2 vs T3 to T8, Table 3 and 4).

237 Maize plants only inoculated of *A. brasilense* was either 1.95% IHFE (Exp.1),
238 8.55% 100 – seed weight, 3.4% harvest index and 17% stem diameter (Exp.2) increased
239 compared when maize plants combination inoculated of *A. brasilense* and fertilized with
240 90 g ha⁻¹ of Mo. However, there was a reduction 10% for NUE, 3.8% LCI and 8.9% stem
241 diameter (Exp.1), 5.8% harvest index and 16% LCI (Exp 3) (T3 an T4 vs T5toT8, Table
242 3 and 4).

243 The effect of *A. brasilense* on seed increased 13.4% NUE, 6.1% shoot dry matter
244 (Exp. 1), 13.3% 100 – seed weight, 2.1% LCI (Exp. 2), 4.8% stem diameter, 8.8% shoot
245 dry matter (Exp. 3), when compared the effect *A. brasilense* on leaf without Mo.
246 However, there was a reduction 8.4% harvest index, 2.8% shoot dry matter (Exp. 2),
247 11.9% harvest index (Exp. 3) (T3 vs T4, Table 3 and 4).

248 The effect of Mo increased 4.4% stem diameter (Exp. 1), 17.8% grain yield,
249 10.4% 100 – seed weight, 11% harvest index, 8.1% stem diameter (Exp. 2), 10.3% NUE,
250 6.6% LCI and 2.4% shoot dry matter (Exp. 3) when compared maize plants inoculated *A.*
251 *brasilense*, regardless of the form of application. However, there was a reduction 5.9%
252 for IHFE and 7.7% shoot dry matter (Exp. 1), 21.5% grain yield, 7.2% harvest index,
253 3.88% stem diameter (Exp.3) (T5 vs T6 to T8, Table 3 and 4).

254 The effect of *A. brasilense* on seed increased 8.1% for 100 – seed weigh, 12.4%
255 NUE, 4.7% LCI, 6% IHFE (Exp. 1), 13.4% grain yield, 8.8% shoot dry matter (Exp.2),
256 8.6% harvest index, 7.9% NUE, 3.7% stem diameter (Exp. 3) when compared *A.*

257 *brasilense* regardless of the form of application. However, there was a reduction 3% for
258 shoot dry matter (Exp. 1), 5.6% LCI, 6.4% IHFE (Exp. 2) and 5.8% shoot dry matter
259 (Exp. 3) (T6 vs T7 and T8, Table 3 and 4).

260 The effect of *A. brasilense* on leaf increased 5.7% for LCI (Exp. 1), 22.5% grain
261 yield, 7.9% harvest index, 20.1% stem diameter, 21.4% IHFE, 6.7% shoot dry matter
262 (Exp. 2) when compared *A. brasilense* on seed with Mo. However, there was a reduction
263 10.5% for NUE, 10.2% for shoot dry matter (Exp. 1), 12.4% harvest index and 5.1%
264 IHFE (Exp. 3) (T7 vs T8, Table 3 and 4).

265 4. DISCUSSION

266 We wanted to understand how the combined use of chemical (N and Mo) and
267 biological (*A. brasilense*) inputs affects the yield and N use efficiency in maize plants
268 cultivated in the sub-humid tropical regions of Brazil. Although it is known that
269 microorganisms favor the growth of crops, less is known about the interactions between
270 diazotrophic (*A. brasiliense*), N rates and molybdenum in maize plants and how this
271 influences plant growth promoting properties, yield and N use efficiency.

272 Notably, N is the nutrient that is most demanded by maize plants and directly
273 affects crop development and yield (Galindo et al. 2020). Nitrogen fixation is a key
274 service in the ecology of plants, but it is an energy-expensive process. Therefore,
275 diazotrophs are critical in the nitrogen fixation process, but generally represent a minority
276 among bacterial communities in plants (Somers et al. 2005; Rajendran et al. 2008; Cassán
277 et al. 2009; Hungria et al. 2010; Cassán et al. 2014), and nitrogen fixation is a tightly
278 regulated process that is deactivated when the fixed nitrogen is available.

279 In the present study, exact mechanisms of the effect of *Azospirillum* inoculation,
280 nitrogen and molybdenum on the development of maize plants were not evaluated, but it
281 is very likely that the improvement in grain yield, LCI and NUE (except experiment 2),

282 reflected in the improvement of shoot dry matter, plant height, length of ear height. The
283 due effect of new production technology is associated inoculated *Azospirillum* its known
284 ability to promote plant growth (Hungria et al. 2010; Galindo et al. 2016; Fukami et al.
285 2017; Martins et al. 2018; Salvo et al. 2018). To Molybdenum, for being a constituent of
286 enzymes (nitrate reductase - NR and glutamine synthetase - GS) and to nitrogen, which
287 enables plant development.

288 The growth-promoting mechanisms due to *A. brasilense* may have improved the
289 plants' ability to exploit the soil more efficiently, as indicated in studies using *A.*
290 *brasilense* (Martins et al. 2018; Fukami et al. 2018a; Fukami et al. 2018b, Leite et al.
291 2019, Zeffa et al. 2019). Furthermore, the application of Mo reduces these adverse effects
292 (N health, increased tissue nitrate concentration), plant growth and development, and thus
293 increases crop yield (Kaiser et al. 2005; Kovács et al. 2015). Finally, the important use of
294 nitrogen maize can be enhanced by molybdenum (Mo), as it is a constituent of enzymes
295 that allow the assimilation of this macronutrient via biological fixation by diazotrophs.
296 microorganisms and/or fertilization with N (Imran et al. 2019).

297 We also observed that the effect of *A. brasilense* on seed increased NUE and shoot
298 dry matter (experiment 1 and 2), when compared the effect *A. brasilense* on leaf without
299 Mo. The stimulation of plant root growth by *A. brasilense* induces an increase in the water
300 absorption and nutrient acquisition rates (including N), which clearly improves the
301 assimilation of N in the biomass and, more generally, plant growth. This capacity is
302 mediated by the bacterial colonization of the roots and their ability to produce different
303 phytohormones, mostly during early stages of plant development and absorb root
304 exudates, it is known maize exudates contain carbohydrates, amino acids and organic
305 acids that serve as carbon sources for bacteria in the rhizosphere (Carvalhais et al. 2011;
306 Van Deynze et al. 2018). On the hand, foliar application of *A. brasilense* creates an

307 unknown interaction between the leaf surface of the plant and the microorganism, which
308 needs to be further investigated (Preininger et al. 2018). Fukami et al. (2016) reported
309 that spray inoculation with *A. brasilense* either on leaves or soil can increase plant growth
310 and can replace 25% of N fertilization on maize. In addition, abiotic factors such as
311 humidity and light have a great impact, furthermore biotic factors including herbivores
312 and the competition between colonizing microbes.

313 Nitrogen use efficiency is controlled by a complex set of interactions among
314 genotype, growing environment, and agronomic management (Hirel et al. 2011; Cañas et
315 al. 2010). In the present study, the new production technology increased NUE. According
316 to Cormier et al. (2013), two strategies can be designed to improve the NUE: increasing
317 the yield on a constant N supply and/or maintaining a high yield by reducing the N supply,
318 a role that can be performed by Mo and as a consequence, helping to achieve a higher
319 grain yield (Munareto, 2016; Galindo et al., 2019a). On the other hand, Calonego et al.
320 (2010) discovered that the absence of Mo foliar supply made for the accumulation of
321 nitrate in common bean leaves: this as a result of the increased nitrogen availability in the
322 soil, which indicated the inefficiency of nitrogen assimilation of plants in the absence of
323 Mo. Srivastava (1997) came to a similar conclusion, stating that in molybdenum-deficient
324 plants, nitrate-reductase activity is often reduced, which results in the buildup of a high
325 concentration of NO_3^- . Mo deficiency likely resulted in an unbalanced nitrogen
326 metabolism, so nitrogen metabolism is seen as a cycle that can be affected by a plant's
327 Mo status (Hu et al., 2002, Yu et al., 2006). Nitrate accumulation in crop plants due to
328 molybdenum deficiency might have serious consequences for human health. Excess
329 nitrate consumption can increase the risk of cancer in adults and causes serious health
330 damage especially in children. (Sanchez-Echaniz et al., 2001)

331 Due to the several mechanisms reported to promote plant growth, Bashan and de-
332 Bashan (2010) proposed the “theory of multiple mechanisms” in which the bacterium
333 leads to a cumulative or sequential pattern of effects, resulting from mechanisms
334 occurring simultaneously or consecutively. According to these authors, there is no single
335 mechanism involved in the promotion of plant growth by specie *A. brasilense* but a
336 combination of a few or many mechanisms determined by inoculant-plant-environment
337 interactions. (Galindo et al. 2020).

338 5. CONCLUSIONS

339 The inoculation with *A. brasilense*, N and Mo, showed potential to be used,
340 because in our study the combination increased 5.1% NUE. However, there was no
341 increase in grain yield. On the other hand, Mo and N increased 17.8% grain yield when
342 compared maize plants inoculated *A. brasilense*.

343 Therefore, studies conducted under conditions that challenge the maize plants in
344 terms of stress, biotic or/and abiotic, are necessary to better understand the role of Mo,
345 applied alone or in combination with growth-promoting bacteria in the sub-humid tropical
346 regions of Brazil

347 Acknowledgements

348 The authors thank the Foundation for the Support of Research and Scientific and
349 Technological Development of Maranhão (FAPEMA) and Coordination for the
350 Improvement of Higher Education Personnel (CAPES) for supporting and for granting a
351 research scholarship and Sandwich PhD Programme respectively to Professor Heder
352 Braun, The Nitro 1000 company for providing the product used (Nitro 1000 Gramíneas).

353 **Competing interests:** The authors declare no competing interests.

354 **Credit authorship contribution statement:**

355 Heder Braun: Conceived and designed the experiment, and wrote and reviewed the
356 original draft of the manuscript;

357 Marcelo Marinho Viana: Performed the experiments in field, collected all samples,
358 collected data in field and performed analysis in the laboratory and wrote and reviewed
359 the original draft of the manuscript.

360 **References**

- 361 Ambrosini, A., Passaglia, L. M. P. 2017. Plant growth-promoting bacteria (PGPB):
362 isolation and screening of PGP activities. *Curr. Prot. Plant Biol.* 2, 190–209.
363 <https://doi.org/10.1002/pb.20054>
- 364 Bashan, Y., & de-Bashan, L. E 2010. How the plant growth-promoting
365 bacterium *Azospirillum* promotes plant growth - a critical assessment. *Adv Agron*
366 108:77–136. [https://doi.org/10.1016/S0065-2113\(10\)08002-8](https://doi.org/10.1016/S0065-2113(10)08002-8)
- 367 Bloch, S. E., Clark, R., Gottlieb, S. S., Wood, L. K., Shah, N., Mak, S. M., Lorigan, J. G.,
368 Johnson, J., Davis-Richardson, A. G., Williams, L., McKellar, M., Soriano, D., Petersen,
369 M., Horton, A., Smith, O., Wu, L., Tung, E., Broglie, R., Tamsir, A., & Temme, K. 2020.
370 Biological nitrogen fixation in maize: Optimizing nitrogenase expression in a root-
371 associated diazotroph. *J. Exp. Bot.* <https://doi: 10.1093/jxb/eraa176>
- 372 Calonego, J. C., Ramos-Junior, E. U., Barbosa, R. D., Leite, G. H. P., & Grassi Filho,
373 H., 2010. Nitrogen topdressing fertilization on common bean with leaf spray of
374 molybdenum (in Portuguese). *Rev. Ci. Agron.* 41, 334: 340.
375 <https://doi.org/10.1590/S1806-66902010000300003>
- 376 Cañas, R. A., Amiour, N., Quilleré, I., & Hirel, B. 2010. An integrated statistical
377 analysis of the genetic variability of nitrogen metabolism in the ear of three maize
378 inbred lines (*Zea mays* L.). *J. Exp. Bot.* 62, 2309–2318.
379 <https://doi.org/10.1093/jxb/erq373>
- 380 Cassán, F., Perrig, D., Sgroy, V., Masciarelli, O., Penna, C., & Luna, V. 2009.
381 *Azospirillum brasilense* Az39 and *Bradyrhizobium japonicum* E109, inoculated singly
382 or in combination, promote seed germination and early seedling growth in corn
383 (*Zea mays* L.) and soybean (*Glycine max* L.). *Eur. J. Soil Biol.* 45 (1), 28–35, [http://](http://dx.doi.org/10.1016/j.ejsobi.2008.08.005)
384 dx.doi.org/10.1016/j.ejsobi.2008.08.005.
- 385 Cassán, F., Vanderleyden, J., & Spaepen, S. 2014. Physiological and agronomical
386 aspects of phytohormone production by model plant-growth-promoting rhizobacteria
387 (pgpr) belonging to the genus *Azospirillum*. *J. Plant Growth Regul.*
388 33 (2), 440–459, <http://dx.doi.org/10.1007/s00344-013-9362-4>
- 389 Cassán, F. D., Okon, Y., & Creus, C.M., 2015. Handbook for
390 *Azospirillum*. *Technical issues and Protocols*. Cham: Springer International
391 Publishing. doi: 10.1007/978-3-319-06542-7

- 392 Cassán, B., & Diaz-Zorita, M. 2017. *Azospirillum* sp. in current agriculture: from the
393 laboratory to the field. *Soil Biol. Biochem.* 103, 117–130. [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.soilbio.2016.08.020)
394 [soilbio.2016.08.020](https://doi.org/10.1016/j.soilbio.2016.08.020)
- 395 Cassán, F., Coniglio, A., López, G., Molina, R., Nieva, S., Carlan, C. L. N., Donadio, F.,
396 Torres, D., Rosas, S., Pedrosa, F. O., Souza, E., Diaz-Zorita, M., Bashan, L. & Mora, V.
397 2020. Everything you must know about *Azospirillum* and its impact on agriculture and
398 beyond. *Biol Fertil Soils* 56, 461–479. <https://doi.org/10.1007/s00374-020-01463-y>
- 399 Carvalhais, L. C., Dennis, P. G., Fedoseyenko, D., Hajirezaei, M. R., Borriss, R., & von
400 Wirén, N. 2011. Root exudation of sugars, amino acids, and organic acids by maize as
401 affected by nitrogen, phosphorus, potassium, and iron deficiency. *J. Plant Nutr. Soil Sci.*
402 174 (1), 3–11, <http://dx.doi.org/10.1002/jpln.201000085>.
- 403 Cormier, F., Faure, S., Dubreuil, P., Heumez, E., Beauchêne, K., Lafarge, S., Praud, S.,
404 & Le Gouis, J. 2013. A multi-environmental study of recent breeding progress on
405 nitrogen use efficiency in wheat (*Triticum aestivum* L.). *Theoretical and Applied*
406 *Genetics*. 2013, 126:3035–3048. doi: 10.1007/s00122-013-2191-9
- 407 Dahal, B., NandaKafle, G., Perkins, L., and Brozel, V. S. 2017. Diversity of free-living
408 nitrogen fixing Streptomyces in soils of the badlands of South Dakota. *Microbiol.*
409 *Res.* 195, 31–39. <https://doi.org/10.1016/j.micres.2016.11.004>
- 410 Fendrik, I., del Gallo, M., Vanderleyden, J., and Zamaroczy, M. 1995.
411 *Azospirillum* VI and Related Microorganisms: Genetics, Physiology, Ecology.
412 Berlin; Heidelberg: Springer-Verlag
- 413 Ferreira, E., Cavalcanti, P. and Nogueira, D. 2014. ExpDes: An R Package for ANOVA
414 and Experimental Designs. *App. Mathe.*, 5, 2952-2958. doi: 10.4236/am.2014.519280.
- 415 Fukami, J., Nogueira, M. A., Araujo, R. S., Hungria, M. 2016. Accessing inoculation
416 methods of maize and wheat with *Azospirillum brasilense*. *AMB Express*. 6, 3–16.
417 <https://doi.org/10.1186/s13568-015-0171-y>
- 418 Fukami, J., Ollero, F. J., Megías, M., and Hungria, M. 2017. Phytohormones
419 and induction of plant-stress tolerance and defense genes by seed and foliar
420 inoculation with *Azospirillum brasilense* cells and metabolites promote maize
421 growth. *AMB Express* 7, 153–163. <https://doi.org/10.1186/s13568-017-0453-7>
- 422 Fukami, J., Abrantes, J. L. F., Cerro, P., Nogueira, M.A., Ollero, F.J., Megías, M., &
423 Hungria, M. 2018a. Revealing strategies of quorum sensing in *Azospirillum brasilense*
424 strains Ab-V5 and Ab-V6. *Arch. Microbiol.* 200, 47–56. [https://doi.org/10.1007/s00203-](https://doi.org/10.1007/s00203-017-1422-x)
425 [017-1422-x](https://doi.org/10.1007/s00203-017-1422-x)
- 426 Fukami, J., Cerezini, P., Hungria, M. 2018b. *Azospirillum*: benefits that go far beyond
427 biological nitrogen fixation. *AMB Express*. 8, 73. [https://doi.org/10.1186/s13568-018-](https://doi.org/10.1186/s13568-018-0608-1)
428 [0608-1](https://doi.org/10.1186/s13568-018-0608-1)
- 429 Galindo, F. S., Teixeira Filho, M. C. M., Buzetti, S., Santini, J. M. K., Alves, C. J.,
430 Nogueira, L. M., Ludkiewicz, M. G. Z., Andreotti, M., Bellotte, J. L. M. 2016. Corn
431 yield and foliar diagnosis affected by nitrogen fertilization and inoculation with
432 *Azospirillum brasilense*. *Rev. Bras. Ciênc. Solo.* 40, e015036. doi:
433 10.1590/18069657rbcsc20150364

- 434 Galindo F. S., Teixeira Filho M. C. M., Buzetti S. S., José, M. K., Ludkiewicz, M. G. Z.,
435 & Baggio, G. 2017. Modes of application of cobalt, molybdenum and *Azospirillum*
436 *brasilense* on soybean yield and profitability. *ver. Bras. de Eng. Agrí. e Amb.*, 21(3), 180-
437 185. <https://doi.org/10.1590/1807-1929/agriambi.v21n3p180-185>
- 438 Galindo, F. S., Rodrigues, W. L., Biagini, A. L. C., Fernandes, G. C., Baratella,
439 E. B. da Silva Junior, C. A., Buzetti, S., & Teixeira Filho, M. C. M. 2019. Assessing
440 forms of application of *Azospirillum brasilense* associated with silicon use on wheat.
441 *Agronomy*, 9(11), 678; <https://doi.org/10.3390/agronomy9110678>
- 442 Galindo, F. S., Pagliari, P. H., Rodrigues, W. L. Pereira, M. R. A., Buzetti, S., Teixeira
443 Filho, M. C. M. 2020. Investigation of *Azospirillum brasilense* inoculation and silicon
444 application on corn yield responses. *J Soil Sci Plant Nutr.* 20, 2406–2418.
445 <https://doi.org/10.1007/s42729-020-00306-x>
- 446 Ganapathy, B. A.; Savalgi, V. P. 2016. Effect of micronutrients on the performance of
447 *Azospirillum brasilense* on the nutrient uptake, growth and yield in maize crop. *Karn.*
448 *Jour. of Agri. Scien.*, 19,66-70
- 449 Gouda, S., Kerry, R. G., Das, G., Paramithiotis, S., Shin, H. S., Patra, J. K. 2018.
450 Revitalization of plant growth promoting rhizobacteria for sustainable development in
451 agriculture. *Micro. Resea.*, 206, 131–140. <https://doi.org/10.1016/j.micres.2017.08.016>
- 452 Kaiser, B. N., Gridley, K. L., Brady, J. N., Phillips, T., & Tyerman, S. D. 2005. The role
453 of molybdenum in agricultural plant production. *Ann Bot* 96:745–754.
454 <https://doi.org/10.1093/aob/mci226>
- 455 Kirkby, E. A.; Römheld, V. 2004. *Micronutrients in Plant Physiology: Functions, Uptake*
456 *and Mobility*. Proceedings 543, The International Fertiliser Society, P. O. Box 4, York,
457 YO32 5YS, Reino Unido.
- 458 Kovács, B., Puskás-Preszner, A., Huzsvai, L., Lévai, L., & Bódi, É. 2015. Effect of
459 molybdenum treatment on molybdenum concentration and nitrate reduction in maize
460 seedlings. *Plant Physiol Biochem* 96:38–44.
461 <https://doi.org/10.1016/j.plaphy.2015.07.013>
- 462 Hirel, B., Tétu, T., Lea, P.J., & Dubois, F. 2011. Improving nitrogen use efficiency in
463 crops for sustainable agriculture. *Sustainability.* 3, 1452–1485.
464 <https://doi.org/10.3390/su3091452>
- 465 Hu, C. X., Wang, Y. H., & Wei, W. X. 2002. Effect of molybdenum applications on
466 concentrations of free amino acids in winter wheat at different growth stages. *J. Plant*
467 *Nutr.*, 25:1487-1499. <https://doi.org/10.1081/PLN-120005404>
- 468 Hungria, M., Campo, R. J., Souza, E. M., Pedrosa, F. O. 2010. Inoculation with selected
469 strains of *Azospirillum brasilense* and *A. lipoferum* improves yields of maize and wheat
470 in Brazil. *Plant Soil*, 331, 413–425. <https://doi.org/10.1007/s11104-009-0262-0>
- 471 Imran. M., Sun. X., Hussain, S., Ali, U., Rana, M. S., Rasul, F., Saleem, M. H., Moussa,
472 M. G., Bhandana, P., Afzal, J. Elyamine, A. M., & Hu, C. X. 2019. Molybdenum-
473 induced effects on nitrogen metabolism enzymes and elemental profile of winter wheat
474 (*Triticum aestivum* L.) under different nitrogen sources. *Int J Mol Sci.* 20:3009.
475 <https://doi.org/10.3390/ijms20123009>

- 476 Leghari S. J., Wahocho N. A., Laghari G. M., Laghari A. H., Bhabhan G. M., Talpur K.
477 A., Bhutto T. A., Wahocho A. S., Lashari A. A. 2016. Role of nitrogen for plant growth
478 and development: A review. *Advances in Environmental Biology*.10(9):209-219.
- 479 Leite, R. C., Santos, J. G. D., Silva, E. L., Alves, C. R. C. R., Hungria, M., Leite, R.C.,
480 & Santos, A.C. 2018. Productivity increase, reduction of nitrogen fertiliser use and
481 drought-stress mitigation by inoculation of Marandu grass (*Urochloa brizantha*) with
482 *Azospirillum brasilense*. *Crop. Pasture Sci.* 70, 61–67. <https://doi.org/10.1071/CP18105>
- 483 Ludueña, L. M., Anzuay, M. S., Angelini, J. G., McIntosh, M., Becker, A., Rupp,
484 O. Goesmann, A., Blom, J., Fabra, A., & Taurian, T. 2018. Strain Serratia sp. S119: a
485 potential biofertilizer for peanut and maize and a model bacterium to study phosphate
486 solubilization mechanisms. *Appl. Soil Ecol.* 126, 107–112.
487 <https://doi.org/10.1016/j.apsoil.2017.12.024>
- 488 Martins, M. R., Jantalia, C. P., Reis, V. M. Döwich, I., Polidoro, J. C., Alves, B. J. R.,
489 Boddey, R. M. & Urquiaga, S. 2018. Impact of plant growth-promoting bacteria on grain
490 yield, protein content, and urea⁻¹⁵ N recovery by maize in a Cerrado Oxisol. *Plant Soil*
491 422, 239–250. <https://doi.org/10.1007/s11104-017-3193-1>
- 492 Moll, R. H., Kamprath, E. J., Jackson, W. A. 1982. Analysis and interpretation of
493 factors which contribute to efficiency of nitrogen utilization 1. *Agronomy journal*,
494 74(3), 562-564. <https://doi.org/10.2134/agronj1982.00021962007400030037x>
- 495 Munareto J.D, Martin T.N, Muller T.M, Nunes U.R, Rosa G.B, Grando L.F.T 2018.
496 Compatibility of *Azospirillum brasilense* with fungicide and insecticide and its effects
497 on the physiological quality of wheat seeds. *Semina: Ci Agrár* 39, 855–864.
498 <http://dx.doi.org/10.5433/1679-0359.2018v39n2p855>
- 499 Noor, M. A. 2017. Nitrogen management and regulation for optimum NUE in maize –
500 mini review. *Cogent Food & Agriculture*, 3(1).
501 <https://doi.org/10.1080/23311932.2017.1348214>
- 502 Oliveira, A. L. M., Santos, O. J. A. P., Marcelino, P. R. F., Milani, K. M. L., Zuluaga, M.
503 Y. A., Zucareli C., & Gonçalves, L. S. A. 2017. Maize Inoculation with *Azospirillum*
504 *brasilense* Ab-V5 Cells Enriched with Exopolysaccharides and Polyhydroxybutyrate
505 Results in High Productivity under Low N Fertilizer Input. *Front. Microbiol.* 8:1873.
506 <https://doi.org/10.3389/fmicb.2017.01873>
- 507 Oliveira, I. J., Fontes, J. R. A., Pereira, B. F. F., & Muniz, A. W. 2018. Inoculation
508 with *Azospirillum* brasiliense increases maize yield. *Chem. Biol. Technol. Agric.*
509 5:6. doi: 10.1186/s40538-018-0118-z
- 510 Pankievicz, V. C. S., Irving, T. B., Maia, L. G. S., & Ané, J. M. 2019. Are we there
511 yet? The long walk towards the development of efficient symbiotic associations
512 between nitrogen-fixing bacteria and non-leguminous crops. *BMC Biol.* 17:99.
513 <https://doi.org/10.1186/s12915-019-0710-0>
- 514 Pereira, N. C. M., Galindo, F. S., Gazola, R. P. D., Dupas, E., Rosa, P. A. L., Mortinho,
515 E. S., Teixeira Filho, M. C. M. 2020. Corn Yield and Phosphorus Use Efficiency
516 Response to Phosphorus Rates Associated with Plant Growth Promoting Bacteria. *Front.*
517 *Environ. Sci.* 8:40. <https://doi.org/10.3389/fenvs.2020.00040>

- 518 Picazevicz, A. A. C., Kusdra, Jorge F., Moreno Andréia de L. 2017. Maize growth in
519 response to *Azospirillum brasilense*, *Rhizobium tropici*, *molybdenum* and
520 nitrogen. *Revista Brasileira de Engenharia Agrícola e Ambiental*, 21(9), 623-
521 627. <https://doi.org/10.1590/1807-1929/agriambi.v21n9p623-627>
- 522 Preininger, C., Sauer, U., Bejarano, A. & Berninger, T. 2018. Concepts and applications
523 of foliar spray for microbial inoculants. *Appl. Microbiol. Biotechnol.* 102, 7265–7282.
524 <https://doi.org/10.1007/s00253-018-9173-4>
- 525 R Core Team. 2019. R: a language and environment for statistical computing. R
526 Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
- 527 RStudio Team. 2019. RStudio: integrated development for R. RStudio, Inc., Boston, MA.
528 <http://www.rstudio.com/>.
- 529 Qi, G., Pan, Z., Sugawa, Y., Andriamanohiarisoamanana, F. J., Yamashiro, T.,
530 Iwasaki, M., Kawamoto, K., Ihara, Ikko & Umetsu, K. 2018. Comparative fertilizer
531 properties of digestates from mesophilic and thermophilic anaerobic digestion of dairy
532 manure: focusing on plant growth promoting bacteria (PGPB) and environmental risk. *J.*
533 *Mat. Cycles Waste Manag.* 20, 1–10. <https://doi.org/10.1007/s10163-018-0708-7>
- 534 Rajendran, G., Sing, F., Desai, A. J., & Archana, G. 2008. Enhanced growth and
535 nodulation of pigeon pea by co-inoculation of *Bacillus* strains with *Rhizobium* spp.
536 *Bioresour. Technol.* 99 (11), 4544–4550, [http://dx.doi.org/10.1016/j.](http://dx.doi.org/10.1016/j.biortech.2007.06.057)
537 [biortech.2007.06.057](http://dx.doi.org/10.1016/j.biortech.2007.06.057)
- 538 Salvo, L.P., Ferrando, L., Fernández-Scavino, A., & Salamone, I. E. G. 2018.
539 Microorganisms reveal what plants do not: Wheat growth and rhizosphere microbial
540 communities after *Azospirillum brasilense* inoculation and nitrogen fertilization under
541 field conditions. *Plant Soil*, 424, 405–417. <https://doi.org/10.1007/s11104-017-3548-7>
- 542 Somers, E., Ptacek, D., Gysegom, P., Srinivasan, M., & Vanderleyden, J. 2005.
543 *Azospirillum brasilense* produces the auxin-like phenylacetic acid by using the
544 key enzyme for indole-3-acetic acid biosynthesis. *Appl. Environ. Microbiol.* 71
545 (4), 1803–1810, <http://dx.doi.org/10.1128/AEM.71.4.1803-1810.2005>
- 546 Srivastava, P. Biochemical significance of molybdenum in crop plants. In *Molybdenum*
547 *in Agriculture*; Gupta, U.C., Ed.; Cambridge University Press: New York, NY, USA,
548 1997; pp. 47–70.
- 549 Soil Survey Staff, 1999. *Soil Taxonomy, a basic classification for making and*
550 *interpreting soil surveys*, 2nd edition Agriculture Handbook 436. USDA, Natural
551 Resources Conservation Service, Washington. 869 p.
- 552 Tedesco, M.J., Gianello, C., Bissani, C., Bohnen, H. and Volkweiss, S.J. 1995. Análise
553 de solo, plantas e outros materiais. [Analysis of soil, plants and other materials.] 2nd
554 Edition, Departamento de Solos da Universidade Federal do Rio Grande do Sul, Porto
555 Alegre, 174.
- 556 Van Deynze, A., Zamora, P., Delaux, P. M., Heitmann, C., Jayaraman, D., Rajasekar,
557 S., Graham, D., Maeda, J., Gibson, D., Schwartz, K. D., Berry, A. M., Bhatnagar, S.,
558 Jospin, G., Darling, A., Jeannotte, R., Lopez, J., Weimer, B. C., Eisen, J. A., Shapiro,
559 H.Y., Ané, J.M., Bennett, A. B. 2018. Nitrogen fixation in a landrace of maize is

- 560 supported by a mucilage-associated diazotrophic microbiota. *PLoS Biol.* 16 (8), 1–21,
561 <http://dx.doi.org/10.1371/journal.pbio.2006352>.
- 562 Valentini, L., F.C. Coelho, and M. dos S. Ferreira. 2005. Leaf nitrogen content and
563 yield of three maize (*Zea mays L.*) cultivars subjected to nitrogen and molybdic
564 fertilization (in Portuguese). *Rev. Ceres.* 52 (302), 567-577.
- 565 Yu, M., Hu, C. X., & Wang, Y. H. 2006. Effects of molybdenum on the intermediates
566 of chlorophyll biosynthesis in winter wheat cultivars under low temperature. *Agric. Sci.*
567 *China.* 5:670-677. [https://doi.org/10.1016/S1671-2927\(06\)60109-0](https://doi.org/10.1016/S1671-2927(06)60109-0)
- 568 Zeffa, D. M., Perini, L. J., Silva, M. B., de Sousa, N. V., Scapim, C. A., Oliveira, A. L.
569 M., Amaral Junior, A. T., & Gonçalves, L. S. A. 2019. *Azospirillum brasilense*
570 promotes increases in growth and nitrogen use efficiency of maize genotypes. *PLoS*
571 *ONE*, 14, e0215332. <https://doi.org/10.1371/journal.pone.0215332>

Table 2. Mean value (\pm SD) of grain yield, 100-seed weight, harvest index, nitrogen use efficiency (NUE) and leaf chlorophyll index (LCI), experiments 1,2 and 3 in the maize under field conditions in research farm, Maranhão State University, Maranhão State, Brazil.

Treatments	Experiment 1				Experiment 2				Experiment 3			
	Grain Yield	100-seed weight	NUE	LCI	Grain Yield	100-seed weight	Harvest Index	LCI	Grain Yield	Harvest Index	NUE	LCI
	t/ha ⁻¹	g	kg kg ⁻¹	-	t/ha ⁻¹	g	%	-	t/ha ⁻¹	%	kg kg ⁻¹	-
1	3,6 \pm 0,2	30,0 \pm 2,1	-	51,5 \pm 1,3	4,3 \pm 0,4	27,3 \pm 1,7	68,4 \pm 1,9	54,0 \pm 3,7	1,8 \pm 0,2	40,9 \pm 1,6	-	73,8 \pm 3,6
2	4,8 \pm 0,7	32,8 \pm 1,2	36,0 \pm 2,2	52,9 \pm 2,6	6,0 \pm 0,4	25,7 \pm 1,5	66,5 \pm 2,5	58,5 \pm 2,6	3,04 \pm 0,7	50,2 \pm 2,9	23,6 \pm 2,1	111,2 \pm 2,8
3	5,2 \pm 0,1	30,9 \pm 1,4	37,1 \pm 1,2	52,5 \pm 2,6	5,8 \pm 0,6	31,3 \pm 1,6	66,6 \pm 2,5	58,6 \pm 2,4	3,2 \pm 0,8	44,6 \pm 3,5	20,7 \pm 2,2	88,9 \pm 2,5
4	4,8 \pm 0,7	32,2 \pm 1,3	32,7 \pm 2,1	52,2 \pm 1,5	5,7 \pm 0,2	27,6 \pm 2,3	72,1 \pm 2,7	62,0 \pm 2,6	2,9 \pm 0,6	50,6 \pm 2,4	22,5 \pm 2,7	88,9 \pm 2,7
5	5,3 \pm 0,5	29,8 \pm 1,2	37,6 \pm 3,7	54,6 \pm 1,2	6,3 \pm 0,6	29,8 \pm 1,6	73,7 \pm 2,4	60,7 \pm 2,1	2,6 \pm 0,7	48,0 \pm 1,5	17,0 \pm 2,2	95,4 \pm 2,9
6	5,8 \pm 0,9	32,6 \pm 1,1	43,8 \pm 2,3	56,6 \pm 1,2	5,6 \pm 0,4	25,6 \pm 1,5	66,7 \pm 1,6	59,8 \pm 1,9	3,6 \pm 0,5	54,6 \pm 2,3	24,2 \pm 1	103,1 \pm 3,7
7	5,3 \pm 0,7	29,2 \pm 1,4	36,4 \pm 3,2	55,5 \pm 2,2	5,5 \pm 0,4	27,2 \pm 2,2	67,7 \pm 1,6	63,9 \pm 2,4	3,2 \pm 0,2	47,0 \pm 2,6	22,6 \pm 1,7	105,6 \pm 2,4
8	5,4 \pm 0,8	30,8 \pm 1,4	40,3 \pm 2,6	52,3 \pm 1,5	4,3 \pm 0,1	27,2 \pm 1,3	62,3 \pm 1,8	62,4 \pm 2,2	2,9 \pm 0,3	52,8 \pm 1,6	20,8 \pm 2,3	102,5 \pm 2,8
Contrast						p value						
1 vs 2 to 8	0,001	0,128	-	0,026	<0,001	0,670	0,720	<0,001	<0,001	<0,001	-	<0,001
2 vs 3 to 8	0,165	0,027	0,195	0,326	0,104	0,014	0,175	0,068	0,910	0,660	0,051	<0,001
3,4 vs 5 to 8	0,140	0,131	<0,001	0,004	0,152	0,013	0,048	0,224	0,993	0,009	0,627	<0,001
3 vs 4	0,419	0,245	0,030	0,810	0,733	0,005	<0,001	0,077	0,583	0,002	0,218	0,974
5 vs 6 to 8	0,529	0,212	0,104	0,908	<0,001	0,004	<0,001	0,378	0,074	0,023	<0,001	<0,001
6 vs 7,8	0,240	0,006	0,003	0,022	0,015	0,143	0,237	0,043	0,116	0,005	0,055	0,519
7 vs 8	0,897	0,136	0,055	0,018	0,001	0,957	0,003	0,400	0,517	0,003	0,223	0,085

Table 3. Mean value (\pm SD) of plant height, stem diameter, insertion height of the first ear (IHFE) and shoot dry matter, experiments 1,2 and 3.

Treatments	Experiment 1			Experiment 2			Experiment 3			
	Stem diameter cm	IHFE	Shoot Dry Matter kg ha ⁻¹	Stem diameter cm	IHFE	Shoot Dry Matter kg ha ⁻¹	Plant height cm	Stem diameter cm	IHFE	Shoot Dry Matter kg ha ⁻¹
1	11,09 \pm 0,3	89,18 \pm 3,6	3575,2 \pm 44,5	11,06 \pm 3,9	46,70 \pm 2,1	3867,08 \pm 78,8	1,11 \pm 0,18	15,1 \pm 0,3	50,9 \pm 4,1	4670,8 \pm 25,8
2	11,23 \pm 0,4	97,60 \pm 1,1	3536,7 \pm 55,1	14,09 \pm 2,8	58,15 \pm 1,9	4361,7 \pm 89,5	1,46 \pm 0,20	17,3 \pm 0,6	54,9 \pm 3,1	4875,6 \pm 20,9
3	10,96 \pm 0,4	95,09 \pm 3,7	3865,11 \pm 59,8	19,32 \pm 2,3	51,32 \pm 3,5	3985,1 \pm 49,9	1,50 \pm 0,10	17,1 \pm 0,2	58,4 \pm 2,5	4658,4 \pm 53,3
4	10,92 \pm 0,5	97,16 \pm 1,2	3644,1 \pm 81,5	18,84 \pm 2,1	53,50 \pm 2,8	4043,7 \pm 72,4	1,41 \pm 0,11	16,3 \pm 0,6	57,5 \pm 3,2	4279,5 \pm 37,7
5	12,63 \pm 0,7	89,47 \pm 2,6	3574,2 \pm 65,9	15,84 \pm 3,6	57,68 \pm 1,2	4100,6 \pm 47,3	1,48 \pm 0,10	16,2 \pm 0,2	58,1 \pm 2,9	4546,6 \pm 44,7
6	12,16 \pm 0,6	98,78 \pm 3,6	3776,3 \pm 70,0	17,68 \pm 2,6	50,81 \pm 2,2	4229,9 \pm 72,5	1,45 \pm 0,06	17,3 \pm 0,7	55,6 \pm 2,2	4270,6 \pm 59,4
7	12,35 \pm 0,5	91,47 \pm 2,3	3700,1 \pm 65,5	18,68 \pm 1,7	60,56 \pm 3,6	3991,5 \pm 77,7	1,47 \pm 0,11	16,4 \pm 0,6	56,1 \pm 2,1	4546,1 \pm 79,6
8	11,72 \pm 0,7	94,12 \pm 2,4	4079,9 \pm 87,7	14,93 \pm 2,2	47,62 \pm 3,3	3723,8 \pm 85,9	1,43 \pm 0,15	16,9 \pm 0,1	58,9 \pm 1,7	4492,4 \pm 72,7
Contrast					p value					
1 vs 2 to 8	0,025	<0,001	0,003	<0,001	<0,001	<0,001	<0,001	<0,001	<0,001	<0,001
2 vs 3 to 8	0,044	0,041	<0,001	0,008	0,003	<0,001	0,901	0,057	0,031	<0,001
3,4 vs 5 to 8	<0,001	0,037	0,377	0,023	0,126	0,181	0,924	0,960	0,427	0,814
3 vs 4	0,913	0,302	<0,001	0,822	0,241	0,043	0,213	0,042	0,544	<0,001
5 vs 6 to 8	0,057	0,003	<0,001	0,338	0,004	0,172	0,577	0,045	0,329	<0,001
6 vs 7,8	0,666	<0,001	0,017	0,525	0,048	<0,001	0,969	0,053	0,154	<0,001
7 vs 8	0,079	0,189	<0,001	0,026	<0,001	<0,001	0,594	0,179	0,066	0,134

Application methods of *Azospirillum brasilense* associated with nitrogen and molybdenum on ecophysiological parameters on maize plants cultivated in the sub-humid tropical regions of Brazil

CAPÍTULO III

1 **Application methods of *Azospirillum brasilense* associated with nitrogen**
2 **and molybdenum on ecophysiological parameters on maize plants**
3 **cultivated in the sub-humid tropical regions of Brazil**

4 **Marcelo Marinho Viana^a, * and Heder Braun^a**

5 ^a*Postgraduate Program in Agroecology, Department of Plant Science and Plant Disease, Maranhão State*
6 *University, São Luís, Maranhão, Brazil.*

7 *** Corresponding author:**

8 *E-mail address: marceloviana.91@gmail.com*

9 **Highlights**

- 10 • The combination of *A. brasilense*, Mo and N increased 12,68% photosynthetic
11 nitrogen use efficiency;
- 12 • Application methods *A. brasilense* on seed average increased 18.5% photosynthetic
13 nitrogen use efficiency;
- 14 • Molybdenum and nitrogen increased at 6.73% photosynthetic nitrogen use efficiency

15

16

17

18

19

20

21

22 ABSTRACT

23 Nitrogen (N) plays a vital role in plant growth, as it is a constituent of nucleotides and proteins.
24 However, the effects induced by combination molybdenum (Mo), N and *Azospirillum*
25 *brasilense* on photosynthetic efficacy have not been investigated. The objective this is study,
26 evaluated the of inoculation methods *A. brasilense*, nitrogen and molybdenum on
27 ecophysiological parameters on maize plants cultivated in the sub-humid tropical regions of
28 Brazil. Three field trials were conducted in a randomized complete block design with four
29 replicates and eight treatment. The treatments were 140 kg ha⁻¹ of N, 90 g ha⁻¹ of Mo and
30 inoculation methods of *A. brasilense* (seed and leaf). In experiment 1, treatments 20 days after
31 top dressing did affect stomatal conductance (g_s), intercellular CO₂ concentration (Ci), leaf
32 chlorophyll index (LCI) and photosynthetic nitrogen use efficiency (PNUE). In experiment 2,
33 treatments did affect Ci, LCI and PNUE 20 and 40 days after top dressing. In experiment 3,
34 treatments did affect A, g_s , 20 days after top dressing, Ci, LCI, PNUE. The average of the
35 experimental results showed a positive effect of the application N, Mo and *A. brasilense*
36 compared to only N or without N. This combination increased PNUE 12,68%. Therefore, it is
37 evident from this study that the application of *A. brasilense* in combination with N and Mo
38 improves N retention, N uptake, photosynthetic apparatus and PNUE of corn grown in sandy
39 soils with poor physical and chemical properties.

40
41 **Keywords:** *Zea mays*; photosynthesis; ecophysiological; biological nitrogen fixation,
42 nitrogen fertilization.

43

44 1. INTRODUCTION

45 Maize (*Zea mays* L.) is a cereal widely cultivated throughout the world due to the
46 important role it plays in the different world agribusiness supply chain. The culture stands out
47 as a source of products for human and animal nutrition (Santos et al, 2017). Nitrogen (N) is

48 typically a limiting nutrient for the production of most of the cereals, and most importantly for
49 maize (Yue et al. 2021), because it is an essential component of all proteins and enzymes,
50 nucleic acids that make up DNA, and chlorophyll that enables the process of photosynthesis in
51 plants (Leghari et al. 2016), moreover nitrogen (N) plays a vital role in plant growth and
52 productivity.

53 N supply has a significant impact on photosynthesis by affecting leaf structure and
54 nitrogen distribution in the leaf (Mu and Chen, 2021). Photosynthesis, or the conversion of
55 light energy into chemical energy, control a variety of physiological, biochemical and
56 molecular processes that substantially contribute to the plant growth and development. (Sener
57 et al. 2016, Imran et al. 2019).

58 Physiologically, the relationship between leaf N and net photosynthetic rate is well
59 documented (Chen et al, 2016). However poor N management has contributed to increased N
60 losses by ammonia (NH_3) volatilization, nitrate (NO_3) leaching and nitrous oxide (N_2O)
61 emissions, which have both economic and environmental consequences (Linguist et al. 2013;
62 Abalos et al. 2014; Martins et al. 2015, 2017; Galindo et al. 2020). These problems indicate
63 the need for a new strategy or alternative, to increased production on maize, nitrogen use
64 efficiency and photosynthetic capacity per unit leaf N.

65 Many studies have shown molybdenum (Mo) plays an essential role in many
66 biochemical processes in plants and is a constituent of nitrogenase (Fageria et al., 2011;
67 Marschner, 2012; Kovács et al, 2015; Barbosa et al, 2021), the enzyme that catalyzes the
68 reduction of atmospheric nitrogen into ammonia, and a cofactor in nitrate reductase (Silva et
69 a., 2017). Mo is less available to plants in acidic soils (typical tropical soils). The limitations
70 of molybdenum in acidic soils indicate the need for strategy Mo foliar supply for plants
71 (Calonego et al., 2010, Kovács et al., 2015).

72 The use of biological techniques such as plant growth-promoting bacteria (PGPB) can
73 represent a sustainable alternative for cereal growth in sub-humid tropical regions of Brazil.
74 (Martins et al., 2018; Galindo et al., 2019). In this sense, inoculation and application of PGPB
75 especially *A. brasilense* is an important strategy in maize cultivation. Several benefits have
76 been attributed to the inoculation with *A. brasilense*, including the supply of N by the biological
77 nitrogen fixation (BNF) process (Hungria et al., 2010; Fukami et al., 2017), stimulation of root
78 growth (Lin et al., 2012; Santi et al., 2013), phosphate solubilization (Rodriguez et al., 2004),
79 and increased tolerance to abiotic (Bulgarelli et al., 2013, Fukami et al., 2018) and biotic
80 (Correa et al., 2008) stresses. In the case of the Brazilian commercial strains of *A. brasilense*
81 Ab-V5 and Ab-V6, the main effects have been attributed to the production of phytohormones
82 (Hungria et al., 2010; Fukami et al., 2017).

83 The most frequent technique of inoculation is via seeds, however, another strategy of
84 application of the PGPB is the foliar, where the PGPB are sprayed over the leaves of the
85 cultivation. The foliar application creates an unknown interaction between the plant leaf surface
86 and the microorganism, which needs to be further investigated (Preininger et al., 2018;
87 Efthimiadou et al., 2020). It should also be noted that the efficiency of inoculation via seeds
88 can be affected by the use of chemicals during treatment, such as fungicides and insecticides,
89 which reinforces the importance of researches and analysis to use supplementary forms of
90 inoculation (Boleta et al., 2020).

91 The need for agriculture that maintains ecosystems and biodiversity, the search for
92 solutions that combine increased production and increasingly sustainable agricultural practices
93 is necessary. Some studies analyze crop production, however, the interaction of the
94 ecophysiological system is still not very explored. Plants can look healthy and no productive.
95 Therefore, the objectives of this study were to evaluate the effects of the combined use of
96 chemical (N and Mo) and biological (*A. brasiliense*) factors on physiological parameters on

97 maize plants grown in the sub-humid tropical regions of Brazil. It was hypothesized that co-
98 application N, Mo, and *A. brasilense* would enhance the photosynthetic efficiency of maize
99 plants.


100 2. MATERIALS E METHODS

101 2.1. *Field site description*

102 Three field trials were conducted in the municipality of São Luís (2°30'S, 44°18'W, 24
103 m above sea level), State of Maranhão, Brazil. The firts trial was stablished from February
104 2018 to May 2018 (Experiment 1), second from October 2018 to January 2019 (Experiment 2)
105 and third from March 2019 to July 2019 (Experiment 3). The region has a hot, semi-humid,
106 equatorial climate, with mean annual rainfall of 2,200 mm and two well-defined seasons: a
107 rainy season from January to June, and a dry season with pronounced water deficits from July
108 to December. The climatic conditions during the trials are shown in Figure 1.

109

110



111

112 **Figure 1.** Rainfall (mm) and maximum and minimum temperatures obtained from the data base
 113 National Institute of Meteorology of Brazil (INMET) during the corn cultivation (all seasons)
 114 in the period from January 2018 to July 2019.

115

116 The trials were conducted under a no-tillage system. The field had not been cultivated
 117 with any agricultural crop for at least 10 years (2003-2013). Maize and cowpea (*Vigna*
 118 *unguiculata* L. Walp) had been cultivated in the field after 2013. The remaining straws from
 119 previous maize and cowpea crops were left on the soil surface. The soil in this area is a Typic
 120 Hapludult according to USDA soil taxonomy (Soil Survey Staff, 1999), with pH in CaCl₂
 121 (soil:solution ratio of 1:2.5) = 5.3; organic matter = 5 g dm⁻³; P (resin) = 31 mg dm⁻³; K⁺ = 1.7
 122 mmolc dm⁻³; Ca²⁺ = 24 mmolc dm⁻³; Mg²⁺ = 15 mmolc dm⁻³; H⁺ + Al³⁺ = 19 mmolc dm⁻³; sum
 123 of bases = 40.7 mmolc dm⁻³; cation exchange capacity at pH 7 = 59.7 mmolc dm⁻³; base
 124 saturation = 68%; and coarse sand = 260 g kg⁻¹, fine sand = 640 g kg⁻¹, silt = 20 g kg⁻¹, and
 125 clay = 120 g kg⁻¹ and texture sandy loam.

126 No mechanical soil preparation was carried out before the maize sowing. At 30 days
 127 before sowing in 2017 was applied 1.0 Mg ha⁻¹ of dolomitic limestone (32% CaO, 15% MgO,
 128 and total neutralizing power of 91%) without incorporation.

129 2.2. Experimental design, treatment, and field management

130 The experimental design was a randomized complete block design with four replicates
 131 and eight treatment (Table 1). There were one N rate 140 kg ha⁻¹, one Mo rate 90 g ha⁻¹ and
 132 inoculation methods of *A. brasilense* (seed and leaf). Each plot consisted of four 4 m-long rows
 133 spaced 0.8 m apart. The outer rows of each plot and 0.5 m from each end of the rows were used
 134 as borders. Treatment arrangement of all seasons 2018/2019 with *Azospirillum brasilense*
 135 (seeds and leaf), nitrogen and molybdenum rates.

136 **Table 1.** Treatment arrangement of all seasons 2018/2019 with *Azospirillum brasilense* (seeds
 137 and leaf), nitrogen and molybdenum rates.

Treatments	Nitrogen (kg/ha ⁻¹)		Molybdenum (g/ha ⁻¹)	Azospirillum seeds (ml/ha ⁻¹)	Azospirillum leaf (ml/ha ⁻¹)
	Sowing	Topdressing	Topdressing	Sowing	Topdressing
1	-	-	-	-	-
2	40	100	-	-	-
3	40	100	-	100	-
4	40	100	-	-	200
5	40	100	90	-	-
6	40	100	90	100	-
7	40	100	90	-	200
8	40	100	90	100	200

138
 139 At sowing furrow, 40 kg ha⁻¹ of N (granulated urea), 80 kg ha⁻¹ of P₂O₅ (simple
 140 superphosphate), and 100 kg ha⁻¹ of K₂O (potassium chloride) were applied in all trials. Seeds
 141 of the maize simple hybrid PIONNER 30F35[®] was sown at a density of four seeds per meter
 142 (5 plants m⁻²). Seedling emergence occurred between four and six days after sowing for all
 143 trials. All plots were irrigated after the sowing with 15 mm of water, using a drip tape system
 144 and 2-day intervals, for a good and uniform seedling emergence. Approximately every 3 days

145 without rainfall after seedling emergence, the plots were irrigated with 15 mm of water (2-hour
146 irrigation). The plants were irrigated when there was no or low rainfall in the previous week.
147 When there was a need to irrigate, we use the drip tape with flat emitters inside, 16mm
148 diameter, thickness from 0.6mm, spacing from 20cm.

149 Topdressed N (100 kg ha⁻¹ of N) (granulated urea) was manually applied when the
150 maize plants had six leaves completely expanded (V6) and was evenly applied along the
151 furrows at 10 cm away from the plants. All plots were immediately irrigated after N fertilizer
152 with approximately 15 mm of water to minimize ammonia volatilization. Topdressed Mo (90
153 g ha⁻¹) (ammonium molybdate) applied when the maize plants had six leaves completely
154 expanded (V6) by spraying the solution, using a sprayer equipped with one cone nozzle (XR
155 11002; Teejet®, Wheaton, USA), with a flow rate of 200 L ha⁻¹ of water. A plastic sheet was
156 used to protect adjacent plots from unwanted spray drift.

157 The seeds and leaves of maize plants were inoculated with *Azospirillum brasilense*,
158 using the commercial strains Ab-V5 and Ab-V6 (Nitro 1000 Gramíneas; Nitro 1000, Cascavel,
159 Brazil) at rates of 100 mL and 200 mL of the liquid inoculant per hectare (2×10⁸ CFU [colony
160 forming unity] mL⁻¹), respectively. These strains have not been used in sub-humid tropical
161 regions of the State of Maranhão, Brazil. The seeds were inoculated one hour before sowing
162 the crop, and the leaves were inoculated at V6 stage by spraying the solution, using a sprayer
163 equipped with one cone nozzle (XR 11002; Teejet®, Wheaton, USA), with a flow rate of 200
164 L ha⁻¹ of water. A plastic sheet was used to protect adjacent plots from unwanted spray drift.

165 Deltamethrin (5 g ha⁻¹ active ingredient) was applied to control fall armyworm
166 (*Spodoptera frugiperda*), when needed. Weeds were controlled by manual hoeing until the N
167 topdressing application.

168 2.3. *Measurements collected*

169 The features were evaluated at the beginning of tasseling stage, (approximately 20 days
170 after topdressed) and milk stage (approximately 40 days after topdressed). The photosynthetic
171 CO₂ assimilation (A), stomatal conductance (g_s) and intercellular CO₂ concentration (C_i) was
172 measured with a portable LI-COR gas exchange system LI-6400 (LI-COR, Lincoln, USA),
173 operating as an open system. The measurements were conducted under a clear and sunny day,
174 and with photosynthetically active radiation of 1,500 μmol m⁻² s⁻¹. The average leaf
175 temperature inside the chamber was 33.6±0.2 °C, and a CO₂ concentration of 400±1 μmol CO₂
176 mol s⁻¹. At least two measurements were performed on third leaf above the ear leaf in two
177 plants, and more were performed when the first two values presented high variation. The
178 average of the four readings was taken as a replicate. The leaf chlorophyll index (LCI), was
179 measured indirectly using a portable non-destructive chlorophyll meter SPAD-502 (Minolta
180 Co., Japan).

181 The same leaves that had been used for the photosynthetic CO₂ assimilation
182 measurements were harvested. Subsequently, 20 leaf disks of approximately 1.8 cm² were
183 taken and the remaining leaf tissues were oven-dried at 70 °C to a constant weight to calculate
184 the specific leaf area (leaf area per unit of dry weight). Total N concentration in leaves were
185 analyzed using the Kjeldahl method (Tedesco et al., 1995). In addition, the photosynthetic N
186 use efficiency (PNUE) was calculated by dividing photosynthetic CO₂ assimilation by specific
187 leaf N (N concentration per unit leaf area) (Sinclair; Horie, 1989; ; Rodríguez-López et al.,
188 2014).

189 2.4. *Statistical analyses*

190 All data were initially tested for homogeneity of variance with O'Neill and Mathews
191 test and normality using the Shapiro and Wilk test. When the effect of treatment was significant

192 ($p < 0.05$), a set of seven orthogonal contrasts (below) was analysed. The precise p values of
193 these contrasts were reported. Values were reported as means \pm SD.

194 Orthogonal contrasts:

195 a) T1 vs. T2 to T8 (this contrast check the effect of N deficiency in maize plants)

196 b) T2 vs. T3 to T8 (this contrast check the effect of new production technology)

197 c) T3 and T4 vs T5 to T8 (this contrast check the effect of Azospirillum versus the combination
198 Mo and Azospirillum)

199 d) T3 vs. T4 (this contrast check the effect of Azospirillum on seed versus Azospirillum on leaf,
200 without Mo application)

201 e) T5 vs T6 to T8 (this contrast check the effect of Mo versus Azospirillum, regardless of the
202 form of application)

203 f) T6 vs T7 and T8 (this contrast check the effect of Azospirillum on seed versus Azospirillum,
204 regardless of the form of application)

205 g) T7 vs T8 (this contrast check the effect of Azospirillum on leaf versus Azospirillum on seed
206 with Mo application).

207 The data were analysed using the statistical software R v. 3.6.1 (R Core Team, 2019)
208 with the RStudio Version 1.2.5019 interface (RStudio Team, 2019) and the ExpDes.pt package
209 (Ferreira; Calvacanti and Nogueira, 2014).

210 3. RESULTS

211 In experiment 1, treatments were evaluated at 20 days after topdressing (20 DATD) did
212 affect stomatal conductance (g_s), intercellular CO_2 concentration (C_i), leaf chlorophyll index
213 (LCI) and photosynthetic nitrogen use efficiency (PNUE). Treatments were evaluated at 40
214 days after topdressing (40 DATD) did affect C_i , LCI, PNUE. Treatments did not affect CO_2
215 assimilation (A) ($p = 0.06$, mean = 43.17 ± 3.66), 20 DATD and CO_2 assimilation ($p = 0.93$,
216 mean = 44.67 ± 3.42) and stomatal conductance ($p = 0.62$, mean = 0.34 ± 0.05) 40 DATD).

217 In experiment 2, treatments were evaluated at 20 DATD did affect C_i , LCI and PNUE
218 and treatments were evaluated at 40 DATD did affect PI_{abs} (40 DATD). Treatments did not
219 affect CO_2 assimilation ($p = 0.124$, mean = 51.1 ± 2.62) and stomatal conductance ($p = 0.39$,
220 mean = 0.55 ± 0.13) at 20 DATD. CO_2 assimilation ($p = 0.09$, mean = 26.22 ± 2.81) and
221 stomatal conductance ($p = 0.57$, mean = 0.59 ± 0.14) at 40 DATD.

222 In experiment 3, treatments were evaluated at 20 DATD did affect A , g_s , C_i , LCI, PNUE
223 and treatments were evaluated at 40 DATD did affect PNUE Treatments did not affect CO_2
224 assimilation ($p = 0.06$, mean = 23.3 ± 2.52) and stomatal conductance ($p = 0.65$, mean = $0.6 \pm$
225 0.68) at 40 DATD.

226 About the contrasts, maize plants N deficiency (T1) was either at 20 DATD in
227 experiment 1 (16.4% LCI, 13.7 % PNUE), experiment 2 (16.5% C_i , 14.5% LCI), experiment
228 3 (2% C_i , 26.1% LCI) and at 40 DATD in experiment 1 (18.5% LCI), experiment 2 (7.2% C_i ,
229 8.9% LCI, 9.9 PNUE) and experiment 3 (7.61% C_i , 21.6% LCI) greater compared when maize
230 plants fertilized of nitrogen. However, at 20 DATD in experiment 1 (4.4% C_i), experiment 3
231 (8.6% A , 22.1% g_s) and at 40 DATD experiment 1 (8.3% C_i , 5.3% PNUE), experiment 3 (6.5%
232 PNUE) was increased with N deficiency (T1 vs T2 to T8, Table 2, 3, 4 and 5).

233 Maize plants fertilized with 140 kg ha^{-1} of nitrogen, 90 g ha^{-1} of Mo and inoculated *A.*
234 *brasilense* (seed and leaf) was either at 20 DATD in experiment 1 (18.2% g_s), experiment 2
235 (3.3% LCI) and at 40 DATD in experiment 1 (18.8% C_i , 5.8% LCI), experiment 2 (2% LCI,
236 11.7% PNUE) and experiment 3 (10.2% LCI, 9.6% PNUE) greater compared when maize
237 plants only fertilized with 140 kg ha^{-1} of nitrogen. However, there was a reduction at 20 DATD
238 in experiment 1 (2.6% C_i , 6.9% LCI), experiment 2 (13.3% PNUE), experiment 3 (10.6% C_i ,
239 20.6% LCI) and at 40 DATD in experiment 1 (12.9% PNUE), experiment 2 (3% C_i),
240 experiment 3 (7.3% C_i) (T2 vs T3 to T8, Table 2, 3, 4 and 4).

241 Maize plants only inoculated of *A. brasilense* was either at 20 DATD in experiment 1
242 (3.4% LCI) and at 40 DATD in experiment 1 (9.3 Ci) and experiment 3 (10.2% LCI) greater
243 compared when maize plants combination inoculated of *A. brasilense* and fertilized with 90 g
244 ha⁻¹ of Mo. However, there was a reduction at 20 DATD in experiment 1 (6.6% Ci, 2.7%
245 PNUE), experiment 2 (10.6% Ci), experiment 3 (3.9% Ci, 16.2% LCI) and at 40 DATD in
246 experiment 1 (0.5% PNUE), experiment 2 (0.1% Ci, 8.8% PNUE) (T3 and T4 vs T5 to T8,
247 Table 2, 3, 4 and 5).

248 The effect of *A. brasilense* on seed increased at 20 DATD in experiment 1 (7.9% Ci
249 and 7.8% PNUE), experiment 2 (11.7% Ci and 17% PNUE), experiment 3 (9.7% Ci) and at 40
250 DATD in experiment 1 (30.7% PNUE) respectively, when compared the effect *A. brasilense*
251 on leaf without Mo. However, there was a reduction at 20 DATD in experiment 1 (27.2% Gs),
252 experiment 3 (16.9% PNUE) and 40 DATD in experiment 1 (10.8% LCI), experiment 2 (3.2%
253 Ci, 12.6% PNUE) and experiment 3 (9% PNUE) (T3 vs T4, Table 2, 3, 4 and 5).

254 The effect of Mo increased at 20 DATD in experiment 1 (3.9 Ci, 17.2% PNUE),
255 experiment 3 (7% A, 13% Gs, 6.7% LCI and 4% PNUE) and 40 DATD in experiment 1 (10%
256 Ci, 10.5% PNUE), experiment 2 (3.1% Ci, 8.7% PNUE), experiment 3 (14.3% LCI) when
257 compared maize plants inoculated *A. brasilense*, regardless of the form of application.
258 However, there was a reduction at 20 DATD in experiment 3 (6.1% Ci) and at 40 DATD in
259 experiment 1 (10.6% LCI), experiment 3 (5.1% PNUE) (T5 vs T6T8, Table 2, 3, 3 and 5).

260 The effect of *A. brasilense* on seed increased at 20 DATD in experiment 2 (5.4%
261 PNUE), experiment 3 (3.2% Ci) and 40 DATD in experiment 1 (6.35% Ci and 7.1% PNUE),
262 experiment 2 (2.3% Ci and 4% LCI) when compared *A. brasilense* regardless of the form of
263 application. However, there was a reduction at 20 DATD in experiment 1 (27.2% gs, 46.7%
264 PNUE) and 40 DATD in experiment 2, 3 (12.7% and 14% PNUE) respectively (T6 vs T7 and
265 T8, Table 2, 3, 4 and 5).

266 The effect of *A. brasilense* on leaf increased at 20 DATD in experiment 1 (17.1% gs
267 and 14.6% Ci), experiment 2 (11.3% Ci and 15.6% PNUE), experiment 3 (14.5% gs, 2.4% Ci)
268 and 40 DATD in experiment 1 (19.6% PNUE), experiment 3 (4.3% Ci and 7.4% LCI) when
269 compared *A. brasilense* on seed with Mo. However, there was a reduction at 40 DATD in
270 experiment 1 (7.6% Ci), experiment 2 (3.4% LCI and 16.6% PNUE) (T7 vs T8, Table 2, 3, 4
271 and 5).

272 6. DISCUSSION

273 In the present study, we wanted to understand how the combined use of chemical and
274 biological inputs affects the Photosynthesis, PNUE and efficiency of photochemistry in maize
275 plants cultivated in the sub-humid tropical regions of Brazil. Nitrogen (N) is an essential but
276 generally limiting nutrient for biological systems (Udvardi et al. 2021) Bacteria of the genus
277 *Azospirillum*, native to the soil, and its association with plants has a series of beneficial
278 responses such as increasing the synthesis of photosynthetic pigments (chlorophyll) (Boleta et
279 al., 2020; Bulegon et al., 2017). Molybdenum (Mo) is an essential micronutrient for higher
280 plants and plays an important role in the photosynthetic process due to its major involvement
281 in the chlorophyll biosynthesis pathway (Yu et al., 2006) and in the chloroplast configuration
282 and ultrastructure (Yu et al., 2005).

283 PNUE is a ratio determined simultaneously by numerator (A) and denominator (foliar
284 nitrogen concentration) (Gou et al. 2016). Photosynthesis is an enzyme-mediated process that
285 largely depends on Ribulose-1, 5-bisphosphate carboxylase-oxygenase (Rubisco enzyme).
286 Rubisco enzyme, as the key enzyme that accounts for up to 30% of total leaf nitrogen, is often
287 positively correlated to nitrogen availability and could directly influence photosynthetic
288 capacity (Zhou, 2005; Pooter and Evans, 1998; Sims et al., 1998). In the present study the
289 feature PNUE is affected in all experiments in 20 and 40 DAAT. This suggested that maize
290 plants have a higher light energy convention and electron transport rate. Maize plants invested

291 N to Rubisco. The CO₂ concentration is higher in the vicinity of Rubisco in maize plants. Thus,
292 a lower amount of Rubisco is sufficient to achieve high photosynthetic rate (Seemann et al.,
293 1984; Sage et al., 1987; Mu and Chen, 2021).

294 Nitrogen availability in leaves is strongly positively correlated with photosynthesis,
295 chloroplast structure, and chlorophyll content (Liu et al., 2018). Mo plays a key role in N
296 absorption and assimilation pathways through enzymatic regulation (Campbell, 2001; Schwarz
297 et al., 2009), indicating that Mo application may trigger many nitrogen-dependent
298 physiological, biochemical and molecular processes through effective N acquisition, leading
299 to plant N overall response to availability. Our results indicated that the effect of the new
300 production technology on PNUE at 20 DAAT is not improved when compared with maize
301 plants only fertilized nitrogen. Probably the Mo leaf spray had a low translocation, due to the
302 acidic pH (<5.2) in soil, resulting in low Mo phloem mobility and xylem certainly affected Mo
303 absorption (Valenciano et al., 2011). Campo and Hungria (2002) reported that Mo foliar
304 spraying presents rapid translocation, after application of five days, they observed a higher
305 concentration of molybdenum in the nodules of soybean plants. On the other hand, the effect
306 of combined use of Mo and N leads increase of 10% in PNUE when compared maize plants
307 inoculated *A. brasilense*.

308 Chlorophyll content is strongly correlated with light, PNUE and the final growth and
309 development of vegetative organs (Liu et al., 2018). In addition, chlorophyll a and b play a key
310 role in photosynthetic CO₂ assimilation, participating in the absorption, transfer and conversion
311 of light energy (Sui et al., 2010). In the present study, the leaf chlorophyll index increased in
312 almost all experiments, due to the positive relationship with N and Mo supply. Suggesting that
313 combined application N and Mo induced significant increases in the chlorophyll a and b
314 contents. Irman et al., 2019 indicates that Mo might have an essential role in the chlorophyll
315 biosynthesis. Yu et al. (2006) report that in their essays the chlorophyll biosynthesis process,

316 the transformation of δ -aminolaevulinic acid (ALA) into uroporphyrinogen III (Uro III),
317 intermediates in the chlorophyll biosynthesis pathway, occurs in the chloroplast matrix instead
318 of the membranes thylakoids and in the absence of Mo the transformation process is blocked
319 and results in a decrease in the chlorophyll content in the leaf tissues.

320 Generally, Gs is often used to denote the extent of opening of stomata and Ci to indicate
321 the assimilation ability of mesophyll cells for CO₂ in plants, which are important indices of the
322 photosynthesis of plants and have close relationships with CO₂ assimilation (Farquhar and
323 Sharkey 1982). In the present study, the disparities in results may have been a function of series
324 environmental factors including water and nutrient status, light, CO₂ levels and temperature.

325 According to the set of results, probably the increase in the efficiency of the use of
326 photosynthetic nitrogen (PNUE) induced by Mo and *A. brasilense* occurred with the
327 improvement of the photosynthetic apparatus and also by the better absorption and assimilation
328 of N.

329 7. CONCLUSIONS

330 The average of the experimental results showed a positive effect of the application N,
331 Mo and *A. brasilense*. This combination increased PNUE 12,68%. Therefore, this study
332 showed that the application of *A. brasilense* in combination with N and Mo improves N
333 retention, N uptake, photosynthetic apparatus and PNUE of corn grown in sandy soils with
334 poor physical and chemical properties.

335 However, further studies should be carried out to explore the molecular basis of Mo-
336 induced changes in the photosynthetic apparatus.

337 Acknowledgements

338 The authors thank the Foundation for the Support of Research and Scientific and Technological
339 Development of Maranhão (FAPEMA) and Coordination for the Improvement of Higher
340 Education Personnel (CAPES) for supporting and for granting a research scholarship and

341 Sandwich PhD Programme respectively to Professor Heder Braun, The Nitro 1000 company
342 for providing the product used (Nitro 1000 Gramíneas).

343 **Competing interests:** The authors declare no competing interests.

344 **Credit authorship contribution statement:**

345 Heder Braun: Conceived and designed the experiment, and wrote and reviewed the original
346 draft of the manuscript;

347 Marcelo Marinho Viana: Performed the experiments in field, collected all samples, collected
348 data in field and performed analysis in the laboratory and wrote and reviewed the original
349 draft of the manuscript.

350 **References**

351 Abalos, D., Jeffery, S., Sanz-Cobena, A., Guardia, G., & Vallejo, A. 2014. Meta-analysis of
352 the effect of urease and nitrification inhibitors on crop productivity and nitrogen use efficiency.
353 *Agric Ecosyst Environ.* 189:136–144. <https://doi.org/10.1016/j.agee.2014.03.036>

354 Barbosa, E. P. A., Sodr , D. N., Braun, H., & Vieira, R. F. 2021. Seeds enriched with
355 molybdenum improve cowpea yield in sub-humid tropical regions of Brazil. *Agronomy*
356 *Journal.* 13:1–10. <https://doi.org/10.1002/agj2.20596>

357 Boleta, E. H. M., Galindo, F. S., Jalal, A., Santini, J. M. K., Rodrigues, W. L., Lima B. H., Arf,
358 O., Silva, M. R., Buzetti, S., & Teixeira Filho, M. C. M. 2020. Inoculation with Growth-
359 Promoting Bacteria *Azospirillum brasilense* and Its Effects on productivity and Nutritional
360 Accumulation of Wheat Cultivars. *Front. Sustain. Food Syst.* 4:607262.
361 [doi:10.3389/fsufs.2020.607262](https://doi.org/10.3389/fsufs.2020.607262)

362 Bulgarelli, D., Schlaeppli, K., Spaepen, S., Ver Loren van Themaat, E., Schulze-Lefert, P.
363 Structure and functions of the bacterial microbiota of plants. 2013. *Annu Rev Plant Biol.*
364 64:807-38. [doi: 10.1146/annurev-arplant-050312-120106](https://doi.org/10.1146/annurev-arplant-050312-120106).

365 Bulegon, L. G., Guimaraes, V. F., Klein, J., Battistus, A. G., Inagaki, A. M., Offemann, L. C.,
366 et al. 2017. Enzymatic activity, gas exchange and production of soybean co-inoculated with
367 *Bradyrhizobium japonicum* and *Azospirillum brasilense*. *Aust J. Crop Sci.* 11, 888–896. [doi:](https://doi.org/10.21475/ajcs.17.11.07.pne575)
368 10.21475/ajcs.17.11.07.pne575

369 Campo, R. J., Hungria, M. 2002. Import ncia dos micronutrientes na fixa o biol gica do N₂.
370 *Informa es Agron micas* 98: 6-9.

- 371 Campbell W. H. 2001. Structure and function of eukaryotic NAD(P)H: nitrate reductase.
372 *Cellular and molecular life sciences: CMLS*, 58:2, 194–204.
373 <https://doi.org/10.1007/PL00000847>
- 374 Chen, Y., Wu, D., Mu, X., Xiao, C., Chen, F., Yuan, L. and Mi, G. 2016. Vertical Distribution
375 of Photosynthetic Nitrogen Use Efficiency and Its Response to Nitrogen in Field-Grown Maize.
376 *Crop Science*, 56: 397- 407. <https://doi.org/10.2135/cropsci2015.03.0170>
- 377 Calonego, J. C., Ramos-Junior, E. U., Barbosa, R. D., Leite, G. H. P., & Grassi Filho, H., 2010.
378 Adubação nitrogenada em cobertura no feijoeiro com suplementação de molibdênio via foliar.
379 *Rev. Ci. Agron.* 41, 334: 340. <https://doi.org/10.1590/S1806-66902010000300003>
- 380 Correa, O. S., Romero, A. M., Soria, M. A., & De Estrada, M. *Azospirillum brasilense* plant
381 genotype interactions modify tomato response to bacterial diseases, and root and foliar
382 microbial communities. In: Cassán, F. D., Garcia De Salamone, I. (Ed.) *Azospirillum* sp.: cell
383 physiology, plant interactions and agronomic research in Argentina. Argentina: Asociación
384 Argentina de Microbiología, 2008. p.87-95.
- 385 Efthimiadou, A., Katsenios, N., Chanioti, S., Giannoglou, M., Djordjevic, N., & Katsaros, G.
386 2020. Effect of foliar and soil application of plant growth promoting bacteria on growth,
387 physiology, yield and seed quality of maize under Mediterranean conditions. *Sci*
388 *Rep* 10, 21060. <https://doi.org/10.1038/s41598-020-78034-6>
- 389 Fageria, N. K., Baligar, V. C. & Jones, C. A. *Growth and mineral nutrition of field crops*. 3.ed.
390 Boca Raton: CRC Press, 2011. 560p.
- 391
392 Farquhar, G. D. & Sharkey, T. D. 1982. Stomatal conductance and photosynthesis. *Annu. Rev.*
393 *Plant Physiol.*, 33: 317-345. <https://doi.org/10.1146/annurev.pp.33.060182.001533>
- 394
395 Ferreira, E., Cavalcanti, P. and Nogueira, D. 2014. ExpDes: An R Package for ANOVA and
396 Experimental Designs. *App. Mathe.*, 5, 2952-2958. doi: 10.4236/am.2014.519280.
- 397 Fukami, J., Ollero, E. J., Megías, M., Hungria, M. 2017. Phytohormones and induction of plant-
398 stress tolerance and defense genes by seed and foliar inoculation with *Azospirillum brasilense*
399 cells and metabolites promote maize growth. *AMB*
400 *Express*, 7:1–13. doi: 10.1186/s13568-017-0453-7
- 401 Fukami, J., Cerezini, P., & Hungria, M. 2018. *Azospirillum*: benefits that go far beyond
402 biological nitrogen fixation,” *AMB Express*, 8:1–12. [https://doi.org/10.1186/s13568-018-0608-](https://doi.org/10.1186/s13568-018-0608-1)
403 1
- 404 Galindo, F. S., Teixeira Filho, M. C. M., Buzetti, S., Pagliari, P. H., Santini, J. M. K., Alves,
405 C. J., Megda, M. M., Nogueira, T. A. R., Andreotti, M. & Arf, O. 2019. Maize yield response
406 to nitrogen rates and sources associated with *Azospirillum brasilense*. *Agron. J.* 111:1985–
407 1997. <https://doi.org/10.2134/agronj2018.07.0481>
- 408 Galindo, F. S., Teixeira Filho, M.C.M., Buzetti, S., Pagliari, P. H., & Santini, J. M. K.
409 2020. Can NBPT urease inhibitor in combination with *Azospirillum brasilense* inoculation

- 410 improve wheat development? *Nutr Cycl Agroecosyst* 117, 131–143.
 411 <https://doi.org/10.1007/s10705-020-10061-1>
- 412 Guo, R., Sun, S. & Liu, B. 2016. Difference in leaf water use efficiency/photosynthetic nitrogen
 413 use efficiency of Bt-cotton and its conventional peer. *Sci Rep* 6, 33539.
 414 <https://doi.org/10.1038/srep33539>
- 415 Hungria, M. R. J., Campo, R. J., Souza, E. M. & Pedrosa, F. O. 2010. Inoculation with selected
 416 strains of *Azospirillum brasilense* and *A. lipoferum* improves yields of maize and wheat in
 417 Brazil. *Plant and Soil*, 331: 413–425, 2010. <https://doi.org/10.1007/s11104-009-0262-0>
- 418 Huang, Z. A., Jiang, D. A., Yang, Y., Sun., J. W. & Jin, S. H. 2004. Effects of Nitrogen
 419 Deficiency on Gas Exchange, Chlorophyll Fluorescence, and Antioxidant Enzymes in Leaves
 420 of Rice Plants. *Photosynthetica*. 42,357–364.
 421 <https://doi.org/10.1023/B:PHOT.0000046153.08935.4c>
- 422 Imran, M., Hu, C., Hussain, S., Rana, M. S., Riaz, M., Afzal, J., Aziz, O., Elyamine, A. M.,
 423 Ismael, M. A. F., & Sun, X. 2019. Molybdenum-induced effects on photosynthetic efficacy of
 424 winter wheat (*Triticum aestivum* L.) under different nitrogen sources are associated with
 425 nitrogen assimilation, *Plant Physiology and Biochemistry*, 141:154-163,
 426 <https://doi.org/10.1016/j.plaphy.2019.05.024>.
- 427 Kovács, B., Puskás-Preszner, A., Huzsvai, L., Lévai, L., & Bódi, E. 2015. Effect of
 428 molybdenum treatment on molybdenum concentration and nitrate reduction in maize seedlings.
 429 *Plant Physiology and Biochemistry*, 96: 38-44. <https://doi.org/10.1016/j.plaphy.2015.07.013>.
- 430 Leghari S. J., Wahocho N. A., Laghari G. M., Laghari A. H., Bhabhan G. M., Talpur K. A.,
 431 Bhutto T. A., Wahocho A. S., Lashari A. A. 2016. Role of nitrogen for plant growth and
 432 development: A review. *Advances in Environmental Biology*.10(9):209-219.
- 433 Liu, Z., Gao, J., Gao, F., Liu, P., Zhao, B., Zhang, J. 2018. Photosynthetic Characteristics and
 434 Chloroplast Ultrastructure of Summer Maize Response to Different Nitrogen Supplies.
 435 *Frontiers in Plant Science*. 9:576. <https://www.frontiersin.org/article/10.3389/fpls.2018.00576>
- 436 Lin, L., Li, Z., Hu, C., Zhang, X., Chang, Z., Yang, L., Li, Y., & An, Q. 2012. Plant growth-
 437 promoting nitrogen fixing enterobacteria are in association with sugarcane plants growing in
 438 Guangxi, China. *Microbes and Environments*. 27: 391–398. doi: 10.1264/jsme2.me11275
- 439 Linquist, B. A., Liu, L., van Kessel, C., & van Groenigen, K. J. 2013. Enhanced efficiency
 440 nitrogen fertilizers for rice systems:meta-analysis of yield and nitrogen uptake. *Field Crops*
 441 *Res* 154:246–254. <https://doi.org/10.1016/j.fcr.2013.08.014>
- 442 Martins, M. R., Jantalia, C. P., Polidoro, J. C., Batista, J. N., Alves, B. J. R., Boddey, R. M., &
 443 Urquiaga, S. 2015. Nitrous oxide and ammonia emissions from N fertilization of maize crop
 444 under no-till in a Cerrado soil. *Soil Till Res* 151:75–81.
 445 <https://doi.org/10.1016/j.still.2015.03.004>
- 446 Martins, M. R., Sant’Anna, S. A. C., Zamanc, M., Santos, R. C., Monteiro, R. C., Alves, B. J.
 447 R., Jantalia, C. P., Boddey, R. M., & Urquiaga, S. 2017. Strategies for the use of urease and
 448 nitrification inhibitors with urea: impact on N₂O and NH₃ emissions, fertilizer-15N recovery

- 449 and maize yield in a tropical soil. *Agric Ecosyst Environ* 247:54–62.
450 <https://doi.org/10.1016/j.agee.2017.06.021>
- 451 Martins, M. R., Jantalia, C. P., Reis, V. M. Döwich, I., Polidoro, J. C., Alves, B. J. R., Boddey,
452 R. M. & Urquiaga, S. 2018. Impact of plant growth-promoting bacteria on grain yield, protein
453 content, and urea-¹⁵ N recovery by maize in a Cerrado Oxisol. *Plant Soil* 422, 239–250.
454 <https://doi.org/10.1007/s11104-017-3193-1>
- 455 Marschner, P. *Mineral nutrition of higher plants*. London: Academic Press, 2012. 889p.
456
- 457 Mu, X., Chen, Q., Chen, F., Yuan, L., & Mi, G. 2017. A RNA-Seq Analysis of the Response
458 of Photosynthetic System to Low Nitrogen Supply in Maize Leaf. *International journal of*
459 *molecular sciences*, 18:2624. <https://doi.org/10.3390/ijms18122624>
460
- 461 Mu, I., Chen, Y. 2021. The physiological response of photosynthesis to nitrogen deficiency.
462 *Plant Physiology and Biochemistry*. 158: 76-82. <https://doi.org/10.1016/j.plaphy.2020.11.019>.
463
- 464 Poorter, H., Evans, J. R. 1998. Photosynthetic nitrogen-use efficiency of species that differ
465 inherently in specific leaf area. *Oecologia* 116, 126–137 (1998).
466 <https://doi.org/10.1007/s004420050560>
- 467 Preininger, C., Sauer, U., Bejarano, A., Berninger, T. 2018. Concepts and applications of foliar
468 spray for microbial inoculants. *Appl. Microbiol. Biotechnol.* 102: 7265–7282. doi:
469 10.1007/s00253-018-9173-4
- 470 R Core Team. 2019. R: a language and environment for statistical computing. R Foundation
471 for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
- 472 RStudio Team. 2019. RStudio: integrated development for R. RStudio, Inc., Boston, MA.
473 <http://www.rstudio.com/>.
- 474 Rodríguez, H., Gonzalez, T., Goire, I., Bashan, Y. 2004. Gluconic acid production and
475 phosphate solubilization by the plant growth-promoting bacterium *Azospirillum* spp.
476 *Naturwissenschaften*. 91:552–555. <https://doi.org/10.1007/s00114-004-0566-0>
- 477 Rodríguez-lópez, N. F., Martins, S. C. V., Cavatte, P. C., Silva, P. E. M., Morais, L. E., Pereira,
478 L. F., Reis, J. V., Avila, R. T., Godoy, A. G., Lavinski, A. O., Damatta, F. M. 2014.
479 Morphological and physiological acclimations of coffee seedlings to growth over a range of
480 fixed or changing light supplies. *Envir. and Exper. Botany*, v. 102, p. 1-10, 2014. <https://doi.org/10.1016/j.envexpbot.2014.01.008>
481
- 482 Santi, C., Bogusz, D., & Franche, C. 2013. Biological nitrogen fixation in non-legume plants.
483 *Annals of Botany*, 111:743–767. <https://doi.org/10.1093/aob/mct048>
- 484 Sage, R. F., Percy, R. W., Seemann, J. R. 1987. The Nitrogen Use Efficiency of C3 and C4
485 Plants: III. Leaf Nitrogen Effects on the Activity of Carboxylating Enzymes in *Chenopodium*
486 *album* (L.) and *Amaranthus retroflexus* (L.). *Plant physiology*, 85(2), 355–359.
487 <https://doi.org/10.1104/pp.85.2.355>
488
- 489 Santos, R. J. dos, Bizzarri, J. H., Barbosa, A. P., & Zucareli, C. 2017. Molybdenum application
490 forms associated to topdressing nitrogen fertilization in the production of maize

- 491 crops. *Agronomy Science and Biotechnology*, 3(2), 105.
 492 <https://doi.org/10.33158/ASB.2017v3i2p105>
 493
- 494 Strasser R.J., Tsimilli-Michael M., & Srivastava A. 2004. Analysis of the Chlorophyll a
 495 Fluorescence Transient. In: Papageorgiou G.C., Govindjee (eds) Chlorophyll a Fluorescence.
 496 *Advances in Photosynthesis and Respiration*. 19. [https://doi.org/10.1007/978-1-4020-3218-](https://doi.org/10.1007/978-1-4020-3218-9_12)
 497 9_12
 498
- 499 Seemann, J. R., Badger, M. R., Berry, J. A. 1984. Variations in the specific activity of ribulose-
 500 1, 5-bisphosphate carboxylase between species utilizing differing photosynthetic pathways.
 501 *Plant Physiol.*, 74:791-794. doi: 10.1104/pp.74.4.791
- 502 Sener, M., Strumpfer, J., Singharoy, A., Hunter, C. N., & Schulten, K. 2016. Overall energy
 503 conversion efficiency of a photosynthetic vesicle. *eLife*, 5, e09541.
 504 <https://doi.org/10.7554/eLife.09541>
- 505 Silva, A., Franzini, V. I., Piccolla, C. D., & Muraoka, T. 2017. Molybdenum supply and
 506 biological fixation of nitrogen by two Brazilian common bean cultivars. *Revista Brasileira de*
 507 *Engenharia Agrícola e Ambiental*. 21: 100-105. [http://dx.doi.org/10.1590/1807-](http://dx.doi.org/10.1590/1807-1929/agriambi.v21n2p100-105)
 508 1929/agriambi.v21n2p100-105
- 509 Sims, D. A., Seemann, J. R., Luo, Y. 1998. The significance of differences in the mechanisms
 510 of photosynthetic acclimation to light, nitrogen and CO₂ for return on investment in leaves.
 511 *Functional Ecology* .12:185–194. <https://doi.org/10.1046/j.1365-2435.1998.00194.x>
- 512 Sinclair, T. R., Horie, T. 1989. Leaf nitrogen, photosynthesis, and crop radiation use efficiency:
 513 a review. *Crop Science*. v. 29, p. 90–98.
- 514 Soil Survey Staff, 1999. Soil Taxonomy, a basic classification for making and interpreting soil
 515 surveys, 2nd edition Agriculture Handbook 436. USDA, Natural Resources Conservation
 516 Service, Washington. 869 p.
- 517 Sui, N., Li, M., Li, K. Song, J., & Wang, B. S. 2010. Increase in unsaturated fatty acids in
 518 membrane lipids of Suaeda salsa L. enhances protection of photosystem II under high salinity.
 519 *Photosynthetica* .48, 623–629. <https://doi.org/10.1007/s11099-010-0080-x>
- 520 Schwarz, G., Mendel, R. & Ribbe, M. 2009. Molybdenum cofactors, enzymes and pathways.
 521 *Nature*. 460, 839–847. <https://doi.org/10.1038/nature08302>
- 522 Tedesco, M. J., Gianello, C., Bissani, C., Bohnen, H. and Volkweiss, S. J. 1995. Análise de
 523 solo, plantas e outros materiais. [Analysis of soil, plants and other materials.] 2nd Edition,
 524 Departamento de Solos da Universidade Federal do Rio Grande do Sul, Porto Alegre, 174.
- 525 Tremblay, W., Zoran C. 2012. Sensing crop nitrogen status with fluorescence indicators. A
 526 review. *Agronomy for Sustainable Development*, 32:451-464. Doi:10.1007/s13593-011-0041-
 527 1.
- 528 Udvardi, M., Below, F. E., Castellano, M. J., Eagle, A. J., Giller, K. E., Ladha, J. K., Liu, X.,
 529 Maaz, T. M., Nova-Franco, B., Raghuram, N., Robertson, G.P., Roy, S., Saha, M., Schmidt,
 530 S., Tegeder, M., York, L. M. & Peters. J. W. 2021. A Research Road Map for Responsible Use
 531 of Agricultural Nitrogen. *Front. Sustain. Food Syst.* 5:660155. doi: 10.3389/fsufs.2021.660155

- 532 Valenciano, J. B., Marcelo, V., Miguelez-Frade, M. M. 2011. Effect of different times and
533 techniques of molybdenum application on chickpea (*Cicer arietinum*) growth and yield. *Span*
534 *J Agric Res.* 9: 1271-1278. <https://bit.ly/3fmQ7Ye>
- 535 Xiaohuan, M., Yanling, C. 2021. The physiological response of photosynthesis to nitrogen
536 deficiency, *Plant Physiology and Biochemistry*, 158:76-82,
537 <https://doi.org/10.1016/j.plaphy.2020.11.019>.
- 538 Yue, K., Li, L., Xie, J., Fudjoe, S. K., Zhang, R., Luo, Z., Anwar, S. 2021. Nitrogen Supply
539 Affects Grain Yield by Regulating Antioxidant Enzyme Activity and Photosynthetic Capacity
540 of Maize Plant in the Loess Plateau. *Agronomy*. 11(6):1094.
541 <https://doi.org/10.3390/agronomy11061094>
- 542 Yu M., Hu C. X., Wang Y. H. 2005. Effect of Mo deficiency on the content of chlorophyll and
543 the ultrastructure of chloroplast in winter wheat cultivars. *Journal of Huazhong Agricultural*
544 *University*, 24: 465–469.
- 545 Yu, M., Hu.C. X., Wang, Y. H. 2006. Effects of molybdenum on the intermediates of
546 chlorophyll biosynthesis in winter wheat cultivars under low temperature. *Agric. Sci. China*,
547 5: 670-677. [https://doi.org/10.1016/S1671-2927\(06\)60109-0](https://doi.org/10.1016/S1671-2927(06)60109-0).
- 548 Zhou, Y. Y. 2005. Cultivation techniques for cotton's high yield and high quality. Golden
549 Shield Press.

Table 2. Effect of treatments evaluated at 20 days after topdressing, mean value (\pm SD) of CO₂ assimilation (A, $\mu\text{mol CO}_2\cdot\text{m}^{-2}\cdot\text{s}^{-1}$), stomatal conductance (g_s $\text{mol m}^{-2}\cdot\text{s}^{-1}$), intercellular CO₂ concentration (Ci, $\mu\text{mol CO}_2/\text{mol}^{-1}$) and leaf chlorophyll index (LCI). Experiments 1,2 and 3.

Treatments	Experiment 1			Experiment 2		Experiment 3			
	g_s	Ci	LCI	Ci	LCI	A	g_s	Ci	LCI
1	0,42 \pm 0,07	143,9 \pm 2,6	41,4 \pm 1,9	105,0 \pm 3,0	51,1 \pm 2,4	40,2 \pm 0,9	0,67 \pm 0,04	151,9 \pm 4,4	70,2 \pm 3,5
2	0,38 \pm 0,03	147,6 \pm 2,9	51,4 \pm 1,7	126,2 \pm 2,9	57,2 \pm 3,7	36,9 \pm 2,0	0,59 \pm 0,08	168,8 \pm 2,9	112,0 \pm 2,6
3	0,37 \pm 0,06	141,9 \pm 3,3	51,7 \pm 2,1	136,5 \pm 1,4	60,5 \pm 3,3	35,0 \pm 2,3	0,55 \pm 0,05	157,6 \pm 2,5	86,3 \pm 5,5
4	0,51 \pm 0,08	131,4 \pm 2,6	49,7 \pm 1,2	122,2 \pm 5,1	59,3 \pm 3,6	37,2 \pm 1,6	0,55 \pm 0,03	143,6 \pm 2,8	87,4 \pm 3,1
5	0,53 \pm 0,11	150,8 \pm 3,3	48,3 \pm 2,3	120,8 \pm 3,9	59,8 \pm 3,9	39,6 \pm 1,7	0,60 \pm 0,02	147,2 \pm 2,4	106,9 \pm 5,1
6	0,39 \pm 0,05	143,8 \pm 2,8	47,9 \pm 2,6	127,6 \pm 3,1	58,7 \pm 3,7	36,7 \pm 2,0	0,54 \pm 0,02	152,8 \pm 4,5	101,9 \pm 4,7
7	0,55 \pm 0,05	156,3 \pm 2,7	48,1 \pm 2,9	130,7 \pm 3,9	62,1 \pm 3,7	37,8 \pm 1,1	0,55 \pm 0,08	159,7 \pm 4,0	104,1 \pm 4,4
8	0,45 \pm 0,02	133,4 \pm 3,9	49,4 \pm 2,9	115,8 \pm 3,5	61,9 \pm 3,2	36,0 \pm 3,2	0,47 \pm 0,05	155,8 \pm 2,6	93,6 \pm 4,7
Contrast				p value					
1 vs 2 to 8	0,297	<0,001	<0,001	<0,001	<0,001	0,009	<0,001	0,017	<0,001
2 vs 3 to 8	0,013	0,007	0,057	0,748	0,098	0,914	0,136	<0,001	<0,001
3,4 vs 5 to 8	0,160	<0,001	0,017	<0,001	0,653	0,121	0,609	0,003	<0,001
3 vs 4	0,004	<0,001	0,193	<0,001	0,619	0,156	0,947	<0,001	0,683
5 vs 6 to 8	0,106	0,002	0,911	0,075	0,592	0,032	0,019	<0,001	0,002
6 vs 7,8	0,010	0,589	0,512	0,068	0,138	0,859	0,364	<0,001	0,165
7 vs 8	0,043	<0,001	0,354	<0,001	0,936	0,140	0,044	0,023	<0,001

Table 3. Effect of treatments evaluated at 20 days after topdressing, mean value (\pm SD) of photosynthetic nitrogen use efficiency (PNUE, $\mu\text{mol CO}_2 \text{ g}^{-1} \text{ N. s}^{-1}$). Experiments 1,2 and 3.

Treatments	Experiment 1	Experiment 2	Experiment 3
	PNUE	PNUE	PNUE
1	86,6 \pm 2,3	154,4 \pm 3,9	107,3 \pm 4,6
2	100,4 \pm 3,5	171,1 \pm 2,2	108,2 \pm 3,4
3	102,1 \pm 5,1	163,9 \pm 3,8	97,5 \pm 2,6
4	94,7 \pm 3,4	140,2 \pm 4,1	117,4 \pm 1,1
5	116,3 \pm 3,8	151,6 \pm 3,7	110,9 \pm 3,5
6	73,4 \pm 1,8	143,5 \pm 4,5	108,5 \pm 2,8
7	109,6 \pm 4,1	164,8 \pm 4,4	105,2 \pm 5,2
8	106,1 \pm 3,9	138,8 \pm 2,9	105,9 \pm 3,6
Contrast	p value		
1 vs 2 to 8	<0,001	0,604	0,813
2 vs 3 to 8	0,957	<0,001	0,719
3,4 vs 5 to 8	0,090	0,132	0,912
3 vs 4	0,014	<0,001	<0,001
5 vs 6 to 8	<0,001	0,2071	0,022
6 vs 7,8	<0,001	0,0013	0,136
7 vs 8	0,209	<0,001	0,738

Table 4. Effect of treatments evaluated at 40 days after topdressing, mean value (\pm SD) of CO₂ assimilation (A, $\mu\text{mol CO}_2\cdot\text{m}^{-2}\cdot\text{s}^{-1}$), stomatal conductance (g_s , $\text{mol m}^{-2}\cdot\text{s}^{-1}$), intercellular CO₂ concentration (Ci, $\mu\text{mol CO}_2/\text{mol}^{-1}$) and leaf chlorophyll index (LCI). Experiments 1,2 and 3.

Treatments	Experiment 1		Experiment 2		Experiment 3	
	Ci	LCI	Ci	LCI	Ci	LCI
1	98,51 \pm 3,1	41,4 \pm 2,8	230,4 \pm 3,8	97,1 \pm 2,3	229,7 \pm 2,5	64,1 \pm 4,5
2	75,82 \pm 3,7	47,0 \pm 2,3	252,7 \pm 1,9	103,6 \pm 2,9	262,1 \pm 3,5	72,3 \pm 3,0
3	86,12 \pm 4,5	48,5 \pm 3,0	247,8 \pm 1,9	109,7 \pm 3,3	248,4 \pm 2,2	88,1 \pm 2,3
4	87,32 \pm 3,9	54,3 \pm 2,9	255,9 \pm 2,8	107,4 \pm 3,7	248,6 \pm 2,0	86,9 \pm 5,5
5	89,60 \pm 2,9	47,5 \pm 2,6	251,5 \pm 1,7	107,9 \pm 2,2	244,2 \pm 2,2	90,9 \pm 5,7
6	95,00 \pm 3,2	51,9 \pm 3,9	239,8 \pm 3,7	108,8 \pm 3,2	243,5 \pm 3,3	74,8 \pm 2,9
7	105,15 \pm 3,6	53,7 \pm 2,6	244,6 \pm 2,7	102,4 \pm 3,6	252,5 \pm 4,6	82,5 \pm 4,8
8	97,71 \pm 3,6	52,1 \pm 3,1	246,4 \pm 3,1	106,5 \pm 1,7	241,6 \pm 4,9	76,4 \pm 3,1
Contrast			p value			
1 vs 2 to 8	<0,001	<0,001	<0,001	<0,001	<0,001	<0,001
2 vs 3 to 8	<0,001	0,006	<0,001	0,034	<0,001	<0,001
3,4 vs 5 to 8	<0,001	0,952	<0,001	0,102	0,056	0,001
3 vs 4	0,650	0,005	<0,001	0,278	0,951	0,676
5 vs 6 to 8	<0,001	0,003	<0,001	0,249	0,400	<0,001
6 vs 7,8	0,009	0,577	<0,001	0,023	0,111	0,076
7 vs 8	0,009	0,420	0,317	0,060	<0,001	0,047

Table 5. Effect of treatments evaluated at 40 days after topdressing, mean value (\pm SD) of photosynthetic nitrogen use efficiency (PNUE, $\mu\text{mol CO}_2 \text{ g}^{-1} \text{ N. s}^{-1}$). Experiments 1,2 and 3.

Treatments	Experiment 1	Experiment 2	Experiment 3
	PNUE	PNUE	PNUE
1	164,0 \pm 2,8	70,4 \pm 3,3	94,2 \pm 4,6
2	174,2 \pm 4,4	69,1 \pm 4,2	81,6 \pm 2,7
3	175,7 \pm 4,6	68,1 \pm 3,5	84,5 \pm 2,7
4	134,4 \pm 4,3	78,0 \pm 3,9	92,8 \pm 4,6
5	164,4 \pm 3,0	88,7 \pm 3,6	86,8 \pm 2,8
6	154,4 \pm 4,6	74,6 \pm 4,9	99,4 \pm 5,0
7	159,0 \pm 4,9	77,6 \pm 2,7	87,8 \pm 4,7
8	127,8 \pm 5,2	90,6 \pm 1,8	86,6 \pm 3,4
Contrast		p value	
1 vs 2 to 8	<0,001	<0,001	0,015
2 vs 3 to 8	<0,001	<0,001	<0,001
3,4 vs 5 to 8	0,049	<0,001	0,393
3 vs 4	<0,001	0,001	0,008
5 vs 6 to 8	<0,001	0,001	0,073
6 vs 7,8	0,002	<0,001	<0,001
7 vs 8	<0,001	<0,001	0,674

CONSIDERAÇÕES FINAIS

A dinâmica de aperfeiçoamento da produtividade dos cereais avança em direção a utilização combinada de produtos químicos e biológico, principalmente no viés da sustentabilidade com produtos de baixo custo.

Este trabalho abre a possibilidade de utilização de manejo nutricional da cultura do milho com a combinação de nitrogênio, molibdênio e bactérias diazotróficas (*A. brasilense*), gerando conhecimento científico sobre a interação de nutrientes e das bactérias diazotróficas em solo do tropico úmido brasileiro.

A tese abre novas linhas de pesquisas, sendo estas:

1. Teste em diferentes condições de solos do tropico úmido brasileiro;
2. Avaliação molecular da interação entre nitrogênio, molibdênio e bactérias diazotrófica na rizosfera;
3. Avaliação de produtividade, eficiência do uso do nitrogênio e eficiência fotossintética do uso do nitrogênio com a dose ótima de nutrientes e concentração ideal de bactérias diazotróficas para os solos do tropico úmido.

ANEXOS



SCIENTIA HORTICULTURAE

AUTHOR INFORMATION PACK

TABLE OF CONTENTS

• Description	p.1
• Audience	p.2
• Impact Factor	p.2
• Abstracting and Indexing	p.2
• Editorial Board	p.2
• Guide for Authors	p.5



ISSN: 0304-4238

DESCRIPTION

Scientia Horticulturae is an international journal publishing research related to **horticultural crops**. Articles in the journal deal with open or protected production of **vegetables, fruits, edible fungi** and **ornamentals** under temperate, subtropical and tropical conditions. Papers in related areas (biochemistry, micropropagation, soil science, plant breeding, plant physiology, phytopathology, etc.) are considered, if they contain information of direct significance to **horticulture**. Papers on the technical aspects of horticulture (engineering, crop processing, storage, transport etc.) are accepted for publication only if they relate directly to the living product. In the case of plantation crops, those yielding a product that may be used fresh (e.g. tropical vegetables, citrus, bananas, and other fruits) will be considered, while those papers describing the processing of the product (e.g. rubber, tobacco, and quinine) will not. The scope of the journal includes all horticultural crops but does not include speciality crops such as, medicinal crops or forestry crops, such as bamboo. Basic molecular studies without any direct application in horticulture will not be considered for this journal.

Types of paper:

1. Original full papers (regular papers)
2. Review articles (should cover a part of the subject of active current interest)
3. Short Communications
 - 3.1 Report of preliminary results of important research (pilot investigation: e.g. no duplications or with other restrictions)
 - 3.2 Newly developed methodology or modification of existing methodology, possibly description of first test.
 - 3.3 Results of the application of an earlier published research methodology on other crops or under different conditions (fact finding or recipes) that are nevertheless of interest to an international readership. Should include new scientific insight.
4. Book Reviews (included in the journal on a range of relevant books which are not more than 2 years old).

Original papers should report the results of original research and should be of international and not only regional interest. The material should not have been previously published elsewhere, except in a preliminary form. Reviews should cover a part of the subject active current interest. They may be submitted or invited.

Short Communications should be as completely documented, both by reference to the literature and description of the experimental procedures employed, as a regular paper. They should not occupy more than 4 printed pages (about 8 manuscript pages, including figures, etc.).

For consultation or suggestions please contact the [Editors-in-Chief](#).

AUDIENCE

Horticulturists, Plant Breeders, Plant Physiologists.

IMPACT FACTOR

2020: 3.463 © Clarivate Analytics Journal Citation Reports 2021

ABSTRACTING AND INDEXING

Ecological Abstracts
 Engineering Village - GEOBASE
 CAB International
 EMBiology
 Current Contents - Agriculture, Biology & Environmental Sciences
 Pascal Francis
 Elsevier BIOBASE
 Research Alert
 Web of Science
 Science Citation Index
 BIOSIS Citation Index
 Scopus

EDITORIAL BOARD

Editors-in-Chief

G. Colla, University of Tuscia, Viterbo, Italy

W.W. Guo, National Key Laboratory of Crop Genetic Improvement, Wuhan, China

Cell engineering and citrus genetic improvement, Genetic & epigenetic studies of citrus polyploids, Integrative omics approaches to reveal the molecular mechanism of male sterility & seedlessness of citrus somatic cybrid, Citrus molecular breeding

S. Kondo, Chiba University, Chiba, Japan

P. Martínez-Gómez, Center for Edaphology and Applied Biology of the Segura River, Murcia, Spain

Molecular Markers, Fruits Breeding, Fruit Genomics, Fruit Transcriptomics, Vegetable Breeding, Biotic and Abiotic stresses

B. Pennisi, University of Georgia College of Agricultural and Environmental Sciences - Griffin Campus, Griffin, Georgia, United States of America

Associate Editors

S. Garcia Martinez

F. Garcia-Sanchez, Center for Edaphology and Applied Biology of the Segura River Soil and Water Conservation Group of Plant nutrition Department, Murcia, Spain

D-L. Guo, Henan University of Science and Technology, Department of Horticulture, Henan, China

Grape molecular breeding, Fruit ripening, Bioinformatics, Genomics, Epigenetics

M. Kyriacou, Agricultural Research Institute, Nicosia, Cyprus

Horticulture, Vegetable science, Grafting, Microgreens, Fruit and vegetable quality, Ripening physiology, Postharvest physiology, Carbohydrate metabolism, Phytochemicals, Functional compounds.

J. Lee, Chung-Ang University, Seoul, South Korea

G. Manganaris, Cyprus University of Technology, Lemesos, Cyprus

Arboriculture, postharvest physiology and technology, quality evaluation, phytochemical profile, fruit ripening syndrome, physiological disorders

G. Qin, Institute of Botany Chinese Academy of Sciences, Beijing, China

Y. Roupael, University of Napoli Federico II Department of Agriculture, Portici, Italy

Fruit quality of vegetable crops through pre-harvest factors, plant nutrition, water and irrigation management, soilless production of vegetables and ornamentals, role of grafting, beneficial micro-

organisms, plants biostimulants in horticultural plants, novel specialty crops such as microgreens and edible flowers

J. Xu, Huazhong Agricultural University, Wuhan, China

Carotenoids, Flavonoids, Limonoids, Flavor, Volatile, Secondary mechanism, Sensory science

Founding Editor

S.J. Wellensiek

Editorial Advisory Board

Z.L. Bie, Huazhong Agricultural University, Wuhan, China

S. Burnett, The University of Maine, Orono, Maine, United States of America

Floriculture, Greenhouse, Organic, Irrigation, Propagation, Moisture sensor

R. I. Cabrera, Rutgers University, Department of Plant Biology, New Jersey, United States of America

E. Candir, Hatay Mustafa Kemal University, Antakya, Turkey

physiology, storage of fruit and vegetables, modified atmosphere packaging

M. Cardarelli, CREA Horticulture and Nursery Research Centre, Pontecagnano, Italy

Plant tissue culture, Propagation, Ornamental and aromatic plants, Plant physiology, Biostimulants, Seed treatments and germination, Grafting

F. Cheng

M. Dorais, Laval University, Quebec, Quebec, Canada

H. Ezura, University of Tsukuba Graduate School of Life and Environmental Sciences, Tsukuba, Japan

Physiology, genomics, genetics, breeding and biotechnology of Solanaceae and Cucurbitaceae plants

S. Garcia, Miguel Hernández University, Orihuela, Alicante, Spain

Genetics, Plant breeding, Molecular markers, Marker-assisted selection, Traditional cultivars, Tomato, Genetic variability study

R.G. Guevara-González, Autonomous University of Queretaro, Queretaro, Mexico

A. Gunes, Ankara University, Ankara, Turkey

Y.J. Hao, Shandong Agricultural University College of Horticulture Science and Engineering, Tai'an, Shandong, China

Molecular biology and biotechnology of fruit trees

W.B. Herppich, Leibniz Institute for Agricultural Engineering and Bioeconomy, Potsdam, Germany

C. Honda, The University of Tokyo, Tokyo, Japan

Apple, Anthocyanin, Deciduous fruit production, Ethylene, Postharvest

T. Jemric, University Clinical Hospital Center Zagreb, Zagreb, Croatia

Postharvest Field, Postharvest Technology of Fruits, including Storage and Postharvest Treatments, Fruit Growing and Genetic Resources

M. Karlsson, University of Alaska Fairbanks, Fairbanks, Alaska, United States of America

Controlled environment agriculture, Greenhouse production, High tunnels, Environmental plant physiology, Lighting conditions, Light emitting diodes, Hydroponics, Floriculture, Northern production conditions, Season extensions

S. Kawabata

J.D. Klein, Agricultural Research Organization Volcani Center, Bet Dagan, Israel

seed, vegetable and fruit physiology, postharvest physiology, plant growth regulators, fruit tree culture and management

H. Krishna, ICAR Central Institute of Arid Horticulture, Bikaner, India

Physiology of horticultural crops, Micropropagation, Mycorrhizal/ Plant Growth Promoting Rhizobia (PGPR) studies, Nursery management, Cultivation measures, Post-harvest management, Evaluation and morphological characterization of crops, Organic cultivation

M. Kyriacou, Agricultural Research Institute, Nicosia, Cyprus

Horticulture, Vegetable science, Grafting, Microgreens, Fruit and vegetable quality, Ripening physiology, Postharvest physiology, Carbohydrate metabolism, Phytochemicals, Functional compounds.

C. Miranda, Public University of Navarre, Pamplona, Spain

Pome fruits, Stone fruits, Viticulture, Modelling, Fruit set, Forecasting, Climate, Yield performance, Plant genetic resources, Genetic diversity, Characterization, Water management, Fertilization, Precision agriculture

B. Ouyang, Huazhong Agricultural University, Wuhan, China

Abiotic stress in plants, Functional Genomics

G. Paliyath, University of Guelph, Guelph, Ontario, Canada

post harvest biology, fruit physiology, biochemistry, storage and technology, nutritional components in fruits, vegetables, membrane biology etc.

S.A. Petropoulos, University of Thessaly School of Agricultural Sciences, Volos, Greece

Bioactive compounds, Essential oils, Fruit quality, Greenhouse production, Horticulture, Salinity, Stress physiology, Vegetable quality, Vegetable production, Water stress

M. Policarpo

D. Savvas, Agricultural University of Athens, Athens, Greece

Irrigation of greenhouse crops, Greenhouse microclimate, Salinity management and strategies for improving the use of saline water in agriculture, Soilless culture, Nutrition and fertilization of vegetable crops, Implications of groundwater and surface water management on nutrient cycling,

Physical properties of growing media and their impact on irrigation management of crops grown on substrates, Cultural practices in vegetable crops

L. Tian, London Research and Development Centre, London, Ontario, Canada

D.W. Turner, University of Western Australia School of Agriculture and Environment, Perth, Australia

Y. Wang, University of Florida Department of Food Science and Human Nutrition, Gainesville, Florida, United States of America

Flavor chemistry, Plant metabolomics, Phytochemical identification, Health benefits of plants

T. West, North Dakota State University, Fargo, North Dakota, United States of America

J. Wu, New Zealand Institute for Plant and Food Research Ltd, Auckland, New Zealand

GUIDE FOR AUTHORS

Your Paper Your Way

We now differentiate between the requirements for new and revised submissions. You may choose to submit your manuscript as a single Word or PDF file to be used in the refereeing process. Only when your paper is at the revision stage, will you be requested to put your paper in to a 'correct format' for acceptance and provide the items required for the publication of your article.

To find out more, please visit the Preparation section below.

INTRODUCTION

Types of paper

1. Original full papers (Regular Papers)
2. Review articles should cover a part of the subject of active current interest.
3. Short Communications
 - 3.1 Report of preliminary results of important research (pilot investigation; e.g. no duplications or with other restrictions).
 - 3.2 Newly developed methodology or modification of existing methodology, possibly description of first test.
 - 3.3 Results of the application of an earlier published research methodology on other crops or under different conditions (fact finding or recipes) that are nevertheless of interest to an international readership.
4. Book Reviews will be included in the journal on a range of relevant books which are not more than 2 years old.

Original papers should report the results of original research. The material should not have been previously published elsewhere, except in a preliminary form. Reviews should cover a part of the subject active current interest. They may be submitted or invited and the full review should not exceed 10'000 words.

Short Communications should be as completely documented, both by reference to the literature and description of the experimental procedures employed, as a regular paper. They should not occupy more than 4 printed pages (about 8 manuscript pages, including figures, etc.).

For consultation or suggestions please contact the Editors-in-Chief.

Submission checklist

You can use this list to carry out a final check of your submission before you send it to the journal for review. Please check the relevant section in this Guide for Authors for more details.

Ensure that the following items are present:

One author has been designated as the corresponding author with contact details:

- E-mail address
- Full postal address

All necessary files have been uploaded:

Manuscript:

- Include keywords
- All figures (include relevant captions)
- All tables (including titles, description, footnotes)
- Ensure all figure and table citations in the text match the files provided
- Indicate clearly if color should be used for any figures in print

Graphical Abstracts / Highlights files (where applicable)

Supplemental files (where applicable)

Further considerations

- Manuscript has been 'spell checked' and 'grammar checked'
- All references mentioned in the Reference List are cited in the text, and vice versa
- Permission has been obtained for use of copyrighted material from other sources (including the Internet)

- A competing interests statement is provided, even if the authors have no competing interests to declare
- Journal policies detailed in this guide have been reviewed
- Referee suggestions and contact details provided, based on journal requirements

For further information, visit our [Support Center](#).

BEFORE YOU BEGIN

Ethics in publishing

Authorship of the paper (ICMJE)

Authorship should be limited to those who have made a significant contribution to the conception, design, execution, and interpretation of the reported study. All those who have made significant contributions should be listed as co-authors. Where there are others who have participated in certain substantive aspects of the research project, they should be acknowledged or listed as contributors. The corresponding author should ensure that all appropriate co-authors and no inappropriate co-authors are included on the paper, and that all co-authors have seen and approved the final version of the paper and have agreed to its submission for publication.

For more information on Ethics in publishing and Ethical guidelines for journal publication see <https://www.elsevier.com/publishingethics> and <https://www.elsevier.com/ethicalguidelines>.

Declaration of competing interest

All authors must disclose any financial and personal relationships with other people or organizations that could inappropriately influence (bias) their work. Examples of potential conflicts of interest include employment, consultancies, stock ownership, honoraria, paid expert testimony, patent applications/registrations, and grants or other funding. All authors, including those *without* competing interests to declare, should create a declaration of competing interest statement (which, where relevant, may specify they have nothing to declare). To do so, authors should use [this tool](#) and upload to the submission system at the Attach Files step. **Please do not convert the .docx template to another file type. Author signatures are not required.**

Submission declaration and verification

Submission of an article implies that the work described has not been published previously (except in the form of an abstract, a published lecture or academic thesis, see '[Multiple, redundant or concurrent publication](#)' for more information), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. To verify originality, your article may be checked by the originality detection service [Crossref Similarity Check](#).

Preprints

Please note that [preprints](#) can be shared anywhere at any time, in line with Elsevier's [sharing policy](#). Sharing your preprints e.g. on a preprint server will not count as prior publication (see '[Multiple, redundant or concurrent publication](#)' for more information).

Preprint posting on SSRN

In support of [Open Science](#), this journal offers its authors a free preprint posting service. Preprints provide early registration and dissemination of your research, which facilitates early citations and collaboration.

During submission to Editorial Manager, you can choose to release your manuscript publicly as a preprint on the preprint server [SSRN](#) once it enters peer-review with the journal. Your choice will have no effect on the editorial process or outcome with the journal. Please note that the corresponding author is expected to seek approval from all co-authors before agreeing to release the manuscript publicly on SSRN.

You will be notified via email when your preprint is posted online and a Digital Object Identifier (DOI) is assigned. Your preprint will remain globally available free to read whether the journal accepts or rejects your manuscript.

For more information about posting to [SSRN](#), please consult the [SSRN Terms of Use](#) and [FAQs](#).

Use of inclusive language

Inclusive language acknowledges diversity, conveys respect to all people, is sensitive to differences, and promotes equal opportunities. Content should make no assumptions about the beliefs or commitments of any reader; contain nothing which might imply that one individual is superior to another on the grounds of age, gender, race, ethnicity, culture, sexual orientation, disability or health condition; and use inclusive language throughout. Authors should ensure that writing is free from bias, stereotypes, slang, reference to dominant culture and/or cultural assumptions. We advise to seek gender neutrality by using plural nouns ("clinicians, patients/clients") as default/wherever possible to avoid using "he, she," or "he/she." We recommend avoiding the use of descriptors that refer to personal attributes such as age, gender, race, ethnicity, culture, sexual orientation, disability or health condition unless they are relevant and valid. When coding terminology is used, we recommend to avoid offensive or exclusionary terms such as "master", "slave", "blacklist" and "whitelist". We suggest using alternatives that are more appropriate and (self-) explanatory such as "primary", "secondary", "blocklist" and "allowlist". These guidelines are meant as a point of reference to help identify appropriate language but are by no means exhaustive or definitive.

Author contributions

For transparency, we encourage authors to submit an author statement file outlining their individual contributions to the paper using the relevant CRediT roles: Conceptualization; Data curation; Formal analysis; Funding acquisition; Investigation; Methodology; Project administration; Resources; Software; Supervision; Validation; Visualization; Roles/Writing - original draft; Writing - review & editing. Authorship statements should be formatted with the names of authors first and CRediT role(s) following. [More details and an example](#)

Changes to authorship

Authors are expected to consider carefully the list and order of authors **before** submitting their manuscript and provide the definitive list of authors at the time of the original submission. Any addition, deletion or rearrangement of author names in the authorship list should be made only **before** the manuscript has been accepted and only if approved by the journal Editor. To request such a change, the Editor must receive the following from the **corresponding author**: (a) the reason for the change in author list and (b) written confirmation (e-mail, letter) from all authors that they agree with the addition, removal or rearrangement. In the case of addition or removal of authors, this includes confirmation from the author being added or removed.

Only in exceptional circumstances will the Editor consider the addition, deletion or rearrangement of authors **after** the manuscript has been accepted. While the Editor considers the request, publication of the manuscript will be suspended. If the manuscript has already been published in an online issue, any requests approved by the Editor will result in a corrigendum.

Article transfer service

This journal is part of our Article Transfer Service. This means that if the Editor feels your article is more suitable in one of our other participating journals, then you may be asked to consider transferring the article to one of those. If you agree, your article will be transferred automatically on your behalf with no need to reformat. Please note that your article will be reviewed again by the new journal. [More information.](#)

Copyright

Upon acceptance of an article, authors will be asked to complete a 'Journal Publishing Agreement' (see [more information](#) on this). An e-mail will be sent to the corresponding author confirming receipt of the manuscript together with a 'Journal Publishing Agreement' form or a link to the online version of this agreement.

Subscribers may reproduce tables of contents or prepare lists of articles including abstracts for internal circulation within their institutions. [Permission](#) of the Publisher is required for resale or distribution outside the institution and for all other derivative works, including compilations and translations. If excerpts from other copyrighted works are included, the author(s) must obtain written permission from the copyright owners and credit the source(s) in the article. Elsevier has [preprinted forms](#) for use by authors in these cases.

For gold open access articles: Upon acceptance of an article, authors will be asked to complete a 'License Agreement' ([more information](#)). Permitted third party reuse of gold open access articles is determined by the author's choice of [user license](#).

Author rights

As an author you (or your employer or institution) have certain rights to reuse your work. [More information](#).

Elsevier supports responsible sharing

Find out how you can [share your research](#) published in Elsevier journals.

Role of the funding source

You are requested to identify who provided financial support for the conduct of the research and/or preparation of the article and to briefly describe the role of the sponsor(s), if any, in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the article for publication. If the funding source(s) had no such involvement then this should be stated.

Open access

Please visit our [Open Access page](#) for more information.

Elsevier Researcher Academy

[Researcher Academy](#) is a free e-learning platform designed to support early and mid-career researchers throughout their research journey. The "Learn" environment at Researcher Academy offers several interactive modules, webinars, downloadable guides and resources to guide you through the process of writing for research and going through peer review. Feel free to use these free resources to improve your submission and navigate the publication process with ease.

Language (usage and editing services)

Please write your text in good English (American or British usage is accepted, but not a mixture of these). Authors who feel their English language manuscript may require editing to eliminate possible grammatical or spelling errors and to conform to correct scientific English may wish to use the [English Language Editing service](#) available from Elsevier's Author Services.

Submission

Our online submission system guides you stepwise through the process of entering your article details and uploading your files. The system converts your article files to a single PDF file used in the peer-review process. Editable files (e.g., Word, LaTeX) are required to typeset your article for final publication. All correspondence, including notification of the Editor's decision and requests for revision, is sent by e-mail.

Submit your article

Please submit your article via <https://www.editorialmanager.com/HORTI/default.aspx>

Referees

Suggesting 5 reviewers is required for submission. Please note that only one suggested reviewer can be from your own country, and suggest four potential reviewers from different countries including full contact details and e-mail addresses. Note that the Editor retains the sole right to decide whether or not the suggested reviewers are used.

PREPARATION**Queries**

For questions about the editorial process (including the status of manuscripts under review) or for technical support on submissions, please visit our [Support Center](#).

NEW SUBMISSIONS

Submission to this journal proceeds totally online and you will be guided stepwise through the creation and uploading of your files. The system automatically converts your files to a single PDF file, which is used in the peer-review process.

As part of the Your Paper Your Way service, you may choose to submit your manuscript as a single file to be used in the refereeing process. This can be a PDF file or a Word document, in any format or layout that can be used by referees to evaluate your manuscript. It should contain high enough quality figures for refereeing. If you prefer to do so, you may still provide all or some of the source files at the initial submission. Please note that individual figure files larger than 10 MB must be uploaded separately.

References

There are no strict requirements on reference formatting at submission. References can be in any style or format as long as the style is consistent. Where applicable, author(s) name(s), journal title/book title, chapter title/article title, year of publication, volume number/book chapter and the article number or pagination must be present. Use of DOI is highly encouraged. The reference style used by the journal will be applied to the accepted article by Elsevier at the proof stage. Note that missing data will be highlighted at proof stage for the author to correct.

Formatting requirements

There are no strict formatting requirements but all manuscripts must contain the essential elements needed to convey your manuscript, for example Abstract, Keywords, Introduction, Materials and Methods, Results, Conclusions, Artwork and Tables with Captions.

If your article includes any Videos and/or other Supplementary material, this should be included in your initial submission for peer review purposes.

Divide the article into clearly defined sections.

Please ensure the text of your paper is double-spaced and has consecutive line numbering - this is an essential peer review requirement.

Figures and tables embedded in text

Please ensure the figures and the tables included in the single file are placed next to the relevant text in the manuscript, rather than at the bottom or the top of the file. The corresponding caption should be placed directly below the figure or table.

Peer review

This journal operates a single anonymized review process. All contributions will be initially assessed by the editor for suitability for the journal. Papers deemed suitable are then typically sent to a minimum of two independent expert reviewers to assess the scientific quality of the paper. The Editor is responsible for the final decision regarding acceptance or rejection of articles. The Editor's decision is final. Editors are not involved in decisions about papers which they have written themselves or have been written by family members or colleagues or which relate to products or services in which the editor has an interest. Any such submission is subject to all of the journal's usual procedures, with peer review handled independently of the relevant editor and their research groups. [More information on types of peer review.](#)

REVISED SUBMISSIONS

Use of word processing software

Regardless of the file format of the original submission, at revision you must provide us with an editable file of the entire article. Keep the layout of the text as simple as possible. Most formatting codes will be removed and replaced on processing the article. The electronic text should be prepared in a way very similar to that of conventional manuscripts (see also the [Guide to Publishing with Elsevier](#)). See also the section on Electronic artwork.

To avoid unnecessary errors you are strongly advised to use the 'spell-check' and 'grammar-check' functions of your word processor.

Article structure

Subdivision - numbered sections

Divide your article into clearly defined and numbered sections. Subsections should be numbered 1.1 (then 1.1.1, 1.1.2, ...), 1.2, etc. (the abstract is not included in section numbering). Use this numbering also for internal cross-referencing: do not just refer to 'the text'. Any subsection may be given a brief heading. Each heading should appear on its own separate line.

Introduction

State the objectives of the work and provide an adequate background, avoiding a detailed literature survey or a summary of the results.

Material and methods

Provide sufficient details to allow the work to be reproduced by an independent researcher. Methods that are already published should be summarized, and indicated by a reference. If quoting directly from a previously published method, use quotation marks and also cite the source. Any modifications to existing methods should also be described.

Results

Results should be clear and concise.

Discussion

This should explore the significance of the results of the work, not repeat them. Avoid extensive citations and discussion of published literature. Results and Discussion should be two separate sections.

Conclusions

The main conclusions of the study may be presented in a short Conclusions section, which may stand alone or form a subsection of a Discussion or Results and Discussion section.

Appendices

If there is more than one appendix, they should be identified as A, B, etc. Formulae and equations in appendices should be given separate numbering: Eq. (A.1), Eq. (A.2), etc.; in a subsequent appendix, Eq. (B.1) and so on. Similarly for tables and figures: Table A.1; Fig. A.1, etc.

Essential title page information

- **Title.** Concise and informative. Titles are often used in information-retrieval systems. Avoid abbreviations and formulae where possible.
- **Author names and affiliations.** Please clearly indicate the given name(s) and family name(s) of each author and check that all names are accurately spelled. You can add your name between parentheses in your own script behind the English transliteration. Present the authors' affiliation addresses (where the actual work was done) below the names. Indicate all affiliations with a lower-case superscript letter immediately after the author's name and in front of the appropriate address. Provide the full postal address of each affiliation, including the country name and, if available, the e-mail address of each author.
- **Corresponding author.** Clearly indicate who will handle correspondence at all stages of refereeing and publication, also post-publication. This responsibility includes answering any future queries about Methodology and Materials. **Ensure that the e-mail address is given and that contact details are kept up to date by the corresponding author.**
- **Present/permanent address.** If an author has moved since the work described in the article was done, or was visiting at the time, a 'Present address' (or 'Permanent address') may be indicated as a footnote to that author's name. The address at which the author actually did the work must be retained as the main, affiliation address. Superscript Arabic numerals are used for such footnotes.

Highlights

Highlights are mandatory for this journal as they help increase the discoverability of your article via search engines. They consist of a short collection of bullet points that capture the novel results of your research as well as new methods that were used during the study (if any). Please have a look at the examples here: [example Highlights](#).

Highlights should be submitted in a separate editable file in the online submission system. Please use 'Highlights' in the file name and include 3 to 5 bullet points (maximum 85 characters, including spaces, per bullet point).

Abstract

A concise and factual abstract is required. The abstract should state briefly the purpose of the research, the principal results and major conclusions. An abstract is often presented separately from the article, so it must be able to stand alone. For this reason, References should be avoided, but if essential, then cite the author(s) and year(s). Also, non-standard or uncommon abbreviations should be avoided, but if essential they must be defined at their first mention in the abstract itself.

Graphical abstract

Although a graphical abstract is optional, its use is encouraged as it draws more attention to the online article. The graphical abstract should summarize the contents of the article in a concise, pictorial form designed to capture the attention of a wide readership. Graphical abstracts should be submitted as a separate file in the online submission system. Image size: Please provide an image with a minimum of 531 × 1328 pixels (h × w) or proportionally more. The image should be readable at a size of 5 × 13 cm using a regular screen resolution of 96 dpi. Preferred file types: TIFF, EPS, PDF or MS Office files. You can view [Example Graphical Abstracts](#) on our information site.

Authors can make use of Elsevier's [Illustration Services](#) to ensure the best presentation of their images and in accordance with all technical requirements.

Keywords

Immediately after the abstract, provide a maximum of 6 keywords, using American spelling and avoiding general and plural terms and multiple concepts (avoid, for example, 'and', 'of'). Be sparing with abbreviations: only abbreviations firmly established in the field may be eligible. These keywords will be used for indexing purposes.

Abbreviations

Define abbreviations that are not standard in this field in a footnote to be placed on the first page of the article. Such abbreviations that are unavoidable in the abstract must be defined at their first mention there, as well as in the footnote. Ensure consistency of abbreviations throughout the article.

Acknowledgements

Collate acknowledgements in a separate section at the end of the article before the references and do not, therefore, include them on the title page, as a footnote to the title or otherwise. List here those individuals who provided help during the research (e.g., providing language help, writing assistance or proof reading the article, etc.).

Formatting of funding sources

List funding sources in this standard way to facilitate compliance to funder's requirements:

Funding: This work was supported by the National Institutes of Health [grant numbers xxxx, yyyy]; the Bill & Melinda Gates Foundation, Seattle, WA [grant number zzzz]; and the United States Institutes of Peace [grant number aaaa].

It is not necessary to include detailed descriptions on the program or type of grants and awards. When funding is from a block grant or other resources available to a university, college, or other research institution, submit the name of the institute or organization that provided the funding.

If no funding has been provided for the research, please include the following sentence:

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Nomenclature and units

Follow internationally accepted rules and conventions: use the international system of units (SI). If other units are mentioned, please give their equivalent in SI.

Authors and Editor(s) are, by general agreement, obliged to accept the rules governing biological nomenclature, as laid down in the *International Code of Botanical Nomenclature*, the *International Code of Nomenclature of Bacteria*, and the *International Code of Zoological Nomenclature*.

All botica (crops, plants, insects, birds, mammals, etc.) should be identified by their scientific names when the English term is first used, with the exception of common domestic animals.

All biocides and other organic compounds must be identified by their Geneva names when first used in the text. Active ingredients of all formulations should be likewise identified.

For chemical nomenclature, the conventions of the *International Union of Pure and Applied Chemistry* and the official recommendations of the *IUPAC-IUB Combined Commission on Biochemical Nomenclature* should be followed.

Math formulae

Present simple formulae in the line of normal text where possible. In principle, variables are to be presented in italics.

Number consecutively any equations that have to be displayed separate from the text (if referred to explicitly in the text).

Subscripts and superscripts should be clear.

Greek letters and other non-Roman or handwritten symbols should be explained in the margin where they are first used. Take special care to show clearly the difference between zero (0) and the letter O, and between one (1) and the letter l.

Give the meaning of all symbols immediately after the equation in which they are first used. For simple fractions use the solidus (/) instead of a horizontal line.

Equations should be numbered serially at the right-hand side in parentheses. In general only equations explicitly referred to in the text need be numbered.

The use of fractional powers instead of root signs is recommended. Also powers of e are often more conveniently denoted by exp.

Levels of statistical significance which can be mentioned without further explanation are: * \leq , ** $P < 0.01$ and *** $P < 0.001$.

In chemical formulae, valence of ions should be given as, e.g., Ca^{2+} , not as Ca^{++} . Isotope numbers should precede the symbols, e.g., ^{18}O .

Footnotes

Footnotes should be used sparingly. Number them consecutively throughout the article. Many word processors build footnotes into the text, and this feature may be used. Should this not be the case, indicate the position of footnotes in the text and present the footnotes themselves separately at the end of the article.

Electronic artwork

General points

- Make sure you use uniform lettering and sizing of your original artwork.
- Preferred fonts: Arial (or Helvetica), Times New Roman (or Times), Symbol, Courier.
- Number the illustrations according to their sequence in the text.
- Use a logical naming convention for your artwork files.
- Indicate per figure if it is a single, 1.5 or 2-column fitting image.
- For Word submissions only, you may still provide figures and their captions, and tables within a single file at the revision stage.
- Please note that individual figure files larger than 10 MB must be provided in separate source files.

A detailed [guide on electronic artwork](#) is available.

You are urged to visit this site; some excerpts from the detailed information are given here.

Formats

Regardless of the application used, when your electronic artwork is finalized, please 'save as' or convert the images to one of the following formats (note the resolution requirements for line drawings, halftones, and line/halftone combinations given below):

EPS (or PDF): Vector drawings. Embed the font or save the text as 'graphics'.

TIFF (or JPG): Color or grayscale photographs (halftones): always use a minimum of 300 dpi.

TIFF (or JPG): Bitmapped line drawings: use a minimum of 1000 dpi.

TIFF (or JPG): Combinations bitmapped line/half-tone (color or grayscale): a minimum of 500 dpi is required.

Please do not:

- Supply files that are optimized for screen use (e.g., GIF, BMP, PICT, WPG); the resolution is too low.
- Supply files that are too low in resolution.
- Submit graphics that are disproportionately large for the content.

Color artwork

Please make sure that artwork files are in an acceptable format (TIFF (or JPEG), EPS (or PDF), or MS Office files) and with the correct resolution. If, together with your accepted article, you submit usable color figures then Elsevier will ensure, at no additional charge, that these figures will appear in color online (e.g., ScienceDirect and other sites) regardless of whether or not these illustrations are reproduced in color in the printed version. **For color reproduction in print, you will receive information regarding the costs from Elsevier after receipt of your accepted article.** Please indicate your preference for color: in print or online only. [Further information on the preparation of electronic artwork.](#)

Figure captions

Ensure that each illustration has a caption. A caption should comprise a brief title (**not** on the figure itself) and a description of the illustration. Keep text in the illustrations themselves to a minimum but explain all symbols and abbreviations used.

Tables

Please submit tables as editable text and not as images. Tables can be placed either next to the relevant text in the article, or on separate page(s) at the end. Number tables consecutively in accordance with their appearance in the text and place any table notes below the table body. Be sparing in the use of tables and ensure that the data presented in them do not duplicate results described elsewhere in the article. Please avoid using vertical rules and shading in table cells.

References

Citation in text

Please ensure that every reference cited in the text is also present in the reference list (and vice versa). Any references cited in the abstract must be given in full. Unpublished results and personal communications are not recommended in the reference list, but may be mentioned in the text. If these references are included in the reference list they should follow the standard reference style of the journal and should include a substitution of the publication date with either 'Unpublished results' or 'Personal communication'. Citation of a reference as 'in press' implies that the item has been accepted for publication.

Reference links

Increased discoverability of research and high quality peer review are ensured by online links to the sources cited. In order to allow us to create links to abstracting and indexing services, such as Scopus, CrossRef and PubMed, please ensure that data provided in the references are correct. Please note that incorrect surnames, journal/book titles, publication year and pagination may prevent link creation. When copying references, please be careful as they may already contain errors. Use of the DOI is highly encouraged.

A DOI is guaranteed never to change, so you can use it as a permanent link to any electronic article. An example of a citation using DOI for an article not yet in an issue is: VanDecar J.C., Russo R.M., James D.E., Ambeh W.B., Franke M. (2003). Aseismic continuation of the Lesser Antilles slab beneath northeastern Venezuela. *Journal of Geophysical Research*, <https://doi.org/10.1029/2001JB000884>. Please note the format of such citations should be in the same style as all other references in the paper.

Web references

As a minimum, the full URL should be given and the date when the reference was last accessed. Any further information, if known (DOI, author names, dates, reference to a source publication, etc.), should also be given. Web references can be listed separately (e.g., after the reference list) under a different heading if desired, or can be included in the reference list.

Data references

This journal encourages you to cite underlying or relevant datasets in your manuscript by citing them in your text and including a data reference in your Reference List. Data references should include the following elements: author name(s), dataset title, data repository, version (where available), year, and global persistent identifier. Add [dataset] immediately before the reference so we can properly identify it as a data reference. The [dataset] identifier will not appear in your published article.

References in a special issue

Please ensure that the words 'this issue' are added to any references in the list (and any citations in the text) to other articles in the same Special Issue.

Reference management software

Most Elsevier journals have their reference template available in many of the most popular reference management software products. These include all products that support [Citation Style Language styles](#), such as [Mendeley](#). Using citation plug-ins from these products, authors only need to select the appropriate journal template when preparing their article, after which citations and bibliographies will be automatically formatted in the journal's style. If no template is yet available for this journal, please follow the format of the sample references and citations as shown in this Guide. If you use reference management software, please ensure that you remove all field codes before submitting the electronic manuscript. [More information on how to remove field codes from different reference management software.](#)

Reference formatting

There are no strict requirements on reference formatting at submission. References can be in any style or format as long as the style is consistent. Where applicable, author(s) name(s), journal title/book title, chapter title/article title, year of publication, volume number/book chapter and the article number or pagination must be present. Use of DOI is highly encouraged. The reference style used by

the journal will be applied to the accepted article by Elsevier at the proof stage. Note that missing data will be highlighted at proof stage for the author to correct. If you do wish to format the references yourself they should be arranged according to the following examples:

Reference style

Text: All citations in the text should refer to:

1. *Single author:* the author's name (without initials, unless there is ambiguity) and the year of publication;
2. *Two authors:* both authors' names and the year of publication;
3. *Three or more authors:* first author's name followed by 'et al.' and the year of publication.

Citations may be made directly (or parenthetically). Groups of references can be listed either first alphabetically, then chronologically, or vice versa.

Examples: 'as demonstrated (Allan, 2000a, 2000b, 1999; Allan and Jones, 1999).... Or, as demonstrated (Jones, 1999; Allan, 2000)... Kramer et al. (2010) have recently shown ...'

List: References should be arranged first alphabetically and then further sorted chronologically if necessary. More than one reference from the same author(s) in the same year must be identified by the letters 'a', 'b', 'c', etc., placed after the year of publication.

Examples:

Reference to a journal publication:

Van der Geer, J., Hanraads, J.A.J., Lupton, R.A., 2010. The art of writing a scientific article. *J. Sci. Commun.* 163, 51-59. <https://doi.org/10.1016/j.Sc.2010.00372>.

Reference to a journal publication with an article number:

Van der Geer, J., Hanraads, J.A.J., Lupton, R.A., 2018. The art of writing a scientific article. *Heliyon*. 19, e00205. <https://doi.org/10.1016/j.heliyon.2018.e00205>.

Reference to a book:

Strunk Jr., W., White, E.B., 2000. *The Elements of Style*, fourth ed. Longman, New York.

Reference to a chapter in an edited book:

Mettam, G.R., Adams, L.B., 2009. How to prepare an electronic version of your article, in: Jones, B.S., Smith, R.Z. (Eds.), *Introduction to the Electronic Age*. E-Publishing Inc., New York, pp. 281-304.

Reference to a website:

Cancer Research UK, 1975. Cancer statistics reports for the UK. <http://www.cancerresearchuk.org/aboutcancer/statistics/cancerstatsreport/> (accessed 13 March 2003).

Reference to a dataset:

[dataset] Oguro, M., Imahiro, S., Saito, S., Nakashizuka, T., 2015. Mortality data for Japanese oak wilt disease and surrounding forest compositions. *Mendeley Data*, v1. <https://doi.org/10.17632/xwj98nb39r.1>.

Reference to software:

Coon, E., Berndt, M., Jan, A., Svyatsky, D., Atchley, A., Kikinon, E., Harp, D., Manzini, G., Shelef, E., Lipnikov, K., Garimella, R., Xu, C., Moulton, D., Karra, S., Painter, S., Jafarov, E., & Molins, S., 2020. *Advanced Terrestrial Simulator (ATS) v0.88 (Version 0.88)*. Zenodo. <https://doi.org/10.5281/zenodo.3727209>.

Journal abbreviations source

Journal names should be abbreviated according to the [List of Title Word Abbreviations](#).

Video

Elsevier accepts video material and animation sequences to support and enhance your scientific research. Authors who have video or animation files that they wish to submit with their article are strongly encouraged to include links to these within the body of the article. This can be done in the same way as a figure or table by referring to the video or animation content and noting in the body text where it should be placed. All submitted files should be properly labeled so that they directly relate to the video file's content. In order to ensure that your video or animation material is directly usable, please provide the file in one of our recommended file formats with a preferred maximum size of 150 MB per file, 1 GB in total. Video and animation files supplied will be published online in the electronic version of your article in Elsevier Web products, including [ScienceDirect](#). Please supply 'stills' with your files: you can choose any frame from the video or animation or make a separate image. These will be used instead of standard icons and will personalize the link to your video data. For more detailed instructions please visit our [video instruction pages](#). Note: since video and animation cannot be embedded in the print version of the journal, please provide text for both the electronic and the print version for the portions of the article that refer to this content.

Data visualization

Include interactive data visualizations in your publication and let your readers interact and engage more closely with your research. Follow the instructions [here](#) to find out about available data visualization options and how to include them with your article.

Supplementary material

Supplementary material such as applications, images and sound clips, can be published with your article to enhance it. Submitted supplementary items are published exactly as they are received (Excel or PowerPoint files will appear as such online). Please submit your material together with the article and supply a concise, descriptive caption for each supplementary file. If you wish to make changes to supplementary material during any stage of the process, please make sure to provide an updated file. Do not annotate any corrections on a previous version. Please switch off the 'Track Changes' option in Microsoft Office files as these will appear in the published version.

Research data

This journal encourages and enables you to share data that supports your research publication where appropriate, and enables you to interlink the data with your published articles. Research data refers to the results of observations or experimentation that validate research findings. To facilitate reproducibility and data reuse, this journal also encourages you to share your software, code, models, algorithms, protocols, methods and other useful materials related to the project.

Below are a number of ways in which you can associate data with your article or make a statement about the availability of your data when submitting your manuscript. If you are sharing data in one of these ways, you are encouraged to cite the data in your manuscript and reference list. Please refer to the "References" section for more information about data citation. For more information on depositing, sharing and using research data and other relevant research materials, visit the [research data](#) page.

Data linking

If you have made your research data available in a data repository, you can link your article directly to the dataset. Elsevier collaborates with a number of repositories to link articles on ScienceDirect with relevant repositories, giving readers access to underlying data that gives them a better understanding of the research described.

There are different ways to link your datasets to your article. When available, you can directly link your dataset to your article by providing the relevant information in the submission system. For more information, visit the [database linking page](#).

For [supported data repositories](#) a repository banner will automatically appear next to your published article on ScienceDirect.

In addition, you can link to relevant data or entities through identifiers within the text of your manuscript, using the following format: Database: xxxx (e.g., TAIR: AT1G01020; CCDC: 734053; PDB: 1XFN).

Mendeley Data

This journal supports Mendeley Data, enabling you to deposit any research data (including raw and processed data, video, code, software, algorithms, protocols, and methods) associated with your manuscript in a free-to-use, open access repository. During the submission process, after uploading your manuscript, you will have the opportunity to upload your relevant datasets directly to *Mendeley Data*. The datasets will be listed and directly accessible to readers next to your published article online.

For more information, visit the [Mendeley Data for journals page](#).

Data in Brief

You have the option of converting any or all parts of your supplementary or additional raw data into a data article published in *Data in Brief*. A data article is a new kind of article that ensures that your data are actively reviewed, curated, formatted, indexed, given a DOI and made publicly available to all upon publication (watch this [video](#) describing the benefits of publishing your data in *Data in Brief*). You are encouraged to submit your data article for *Data in Brief* as an additional item directly alongside the revised version of your manuscript. If your research article is accepted, your data article will automatically be transferred over to *Data in Brief* where it will be editorially reviewed, published

open access and linked to your research article on ScienceDirect. Please note an [open access fee](#) is payable for publication in *Data in Brief*. Full details can be found on the [Data in Brief website](#). Please use [this template](#) to write your *Data in Brief* data article.

MethodsX

You have the option of converting relevant protocols and methods into one or multiple MethodsX articles, a new kind of article that describes the details of customized research methods. Many researchers spend a significant amount of time on developing methods to fit their specific needs or setting, but often without getting credit for this part of their work. MethodsX, an open access journal, now publishes this information in order to make it searchable, peer reviewed, citable and reproducible. Authors are encouraged to submit their MethodsX article as an additional item directly alongside the revised version of their manuscript. If your research article is accepted, your methods article will automatically be transferred over to MethodsX where it will be editorially reviewed. Please note an open access fee is payable for publication in MethodsX. Full details can be found on the [MethodsX website](#). Please use [this template](#) to prepare your MethodsX article.

Data statement

To foster transparency, we encourage you to state the availability of your data in your submission. This may be a requirement of your funding body or institution. If your data is unavailable to access or unsuitable to post, you will have the opportunity to indicate why during the submission process, for example by stating that the research data is confidential. The statement will appear with your published article on ScienceDirect. For more information, visit the [Data Statement page](#).

AFTER ACCEPTANCE

Online proof correction

To ensure a fast publication process of the article, we kindly ask authors to provide us with their proof corrections within two days. Corresponding authors will receive an e-mail with a link to our online proofing system, allowing annotation and correction of proofs online. The environment is similar to MS Word: in addition to editing text, you can also comment on figures/tables and answer questions from the Copy Editor. Web-based proofing provides a faster and less error-prone process by allowing you to directly type your corrections, eliminating the potential introduction of errors.

If preferred, you can still choose to annotate and upload your edits on the PDF version. All instructions for proofing will be given in the e-mail we send to authors, including alternative methods to the online version and PDF.

We will do everything possible to get your article published quickly and accurately. Please use this proof only for checking the typesetting, editing, completeness and correctness of the text, tables and figures. Significant changes to the article as accepted for publication will only be considered at this stage with permission from the Editor. It is important to ensure that all corrections are sent back to us in one communication. Please check carefully before replying, as inclusion of any subsequent corrections cannot be guaranteed. Proofreading is solely your responsibility.

Offprints

The corresponding author will, at no cost, receive a customized [Share Link](#) providing 50 days free access to the final published version of the article on [ScienceDirect](#). The Share Link can be used for sharing the article via any communication channel, including email and social media. For an extra charge, paper offprints can be ordered via the offprint order form which is sent once the article is accepted for publication. Both corresponding and co-authors may order offprints at any time via Elsevier's [Author Services](#). Corresponding authors who have published their article gold open access do not receive a Share Link as their final published version of the article is available open access on ScienceDirect and can be shared through the article DOI link.

AUTHOR INQUIRIES

Visit the [Elsevier Support Center](#) to find the answers you need. Here you will find everything from Frequently Asked Questions to ways to get in touch.

You can also [check the status of your submitted article](#) or find out [when your accepted article will be published](#).

© Copyright 2018 Elsevier | <https://www.elsevier.com>